



RESEARCH ARTICLE

Comparison of Estimation Method in Diagnostic Meta-Analysis: An Application in Dentistry

Merve Parmaksiz¹, Hayal Boyacioğlu^{1*} , Pelin Güneri²  and Nezaket Ezgi Özer² 

¹Department of Statistics, Ege University Faculty of Science, Bornova, Izmir, Turkey

²Department of Maxillofacial Radiology, Ege University Faculty of Dentistry, Bornova, Izmir, Turkey

*Corresponding author: Hayal Boyacioğlu, Department of Statistics, Ege University Faculty of Science, Bornova, Izmir, Turkey



Abstract

In this study, the objective was to compare different estimation methods in diagnostic meta-analysis. In this scope, DerSimonian and Laird (DL), Restricted Maximum Likelihood (REML), Sidik and Jonkman (SJ), Hedges and Olkin (HO), Maximum Likelihood (ML), Paule and Mandel (PM) estimation methods were examined. In the implementation part, effectiveness of Clinical Oral Examination (COE) in predicting the diagnosis of histological dysplasia or Oral Squamous Cell Carcinoma (OSCC) was studied. Meta analysis was performed for the data set obtained from 24 studies in accordance with the criteria. Odds Ratio (OR) was used as the effect size. In meta analysis of the random effect model, according to the DerSimonian and Laird (DL) method, the pooled sensitivity value of COE was calculated as 0.953 (95% CI: 0.895-0.979), pooled selectivity was 0.25 (95% CI: 0.124-0.44), and pooled odds ratio was OR = 6.031 (95% CI: 2.208-16.471). According to these results, it can be concluded that COE was not effective in diagnosis. Among the other estimation methods, DerSimonian and Laird (DL) presented the lowest value for I^2 and τ^2 ($I^2 = 66.63\%$, $\tau^2 = 3.489$).

Keywords

Meta-analysis, Diagnostic test, DerSimonian and Laird, Odds ratio

Introduction

In order to acquire trustworthy findings from a scientific research, it is essential to design a comprehensive study plan, to appropriately collect data, to select adequate statistical methods for evaluation and to interpret the results accurately.

Therefore, both the insecurity arising from impractical data, and the inconsistent and contradictory results can be eliminated by combining the previous studies on the same topic. This approach is defined as “meta-analysis” which provides a joint and accurate decision-making opportunity [1]. Meta-analysis aims to predict the related parameters more accurately by increasing the sample size via statistical analysis of the results obtained from the published or unpublished individual studies which are related to a special topic [2,3]. Meta-analysis has been used in 1980s mostly to assess the clinical efficacy of individual medical interventions and since then, it has been a required and advocated statistical analysis in various disciplines [4].

Today, a large number of diseases can be diagnosed and treated. Diagnostic tests which confirm the presence or absence of a disease, give information about the prognosis of the disease and in certain situations, determine the response to treatment have an essential role in medical field [5].

The estimation methods used in meta-analysis have been investigated by numerous studies. Viechtbauer, et al. showed that the Paule and Mandel (PM) estimator is the same as the so-called empirical Bayes estimator [6]. The PM estimation method retains many of the advantages of the method of moments, because it is semiparametric and requires no convergence diagnostics [7]. The moment-based method proposed by DerSimonian and Laird (DL) is most commonly used to estimate the heterogeneity variance. DL method is the

Table 1: 2 × 2 Contingency table.

		References Test		
		With Disease	Without Disease	Total
Test	Positive	a (TP)	b (FP)	a + b
	Negative	c (FN)	d (TN)	c + d
	Total	a + c	b + d	a + b + c + d

Table 2: Diagnostic test indicators.

Test Indicator	Formula
Sensitivity	$Sensitivity = \frac{a}{a + c}$
Specificity	$Specificity = \frac{d}{d + b}$
False Positive Rate	$False\ Positive\ Rate = \frac{b}{b + d}$
False Negative Rate	$False\ Negative\ Rate = \frac{c}{c + a}$
Accuracy	$Accuracy = \frac{a + d}{a + b + c + d}$
PPV (Positive Predictive Value)	$PPV = \frac{a}{a + b}$
NPV (Negative Predictive Value)	$NPV = \frac{d}{c + d}$
PLR (Positive Likelihood Ratio)	$LR+ = \frac{Sensitivity}{1 - Specificity}$
NLR (Negative Likelihood Ratio)	$LR- = \frac{1 - Sensitivity}{Specificity}$
OR (Odds Ratio)	$DOR = \frac{\frac{Sensitivity}{1 - Sensitivity}}{\frac{1 - Specificity}{Specificity}}$
YI (Youden Index)	YI = Sensitivity + Specificity - 1

default approach in many software routines. Simulation studies have found that the method can be biased and thus, other methods have been introduced [3]. The maximum likelihood (ML) method is asymptotically efficient, but requires an iterative solution. The Sidikve Jonkman (SJ) estimator has methodological similarities with the PM estimator. Although the Hedges ve Olkin (HO) estimator is simple to compute and does not require an iterative numerical solution, it is not widely used [8,9]. On the other hand, restricted maximum likelihood (REML) estimation is a generally well-known estimation technique in the statistical literature [10]. Bowden, et al. conducted an empirical study by comparing DL and PM estimation methods and stated that as the variance between studies increased, τ^2 value of PM was greater

than τ^2 obtained from DL [11].

Study Method

Diagnostic test

Diagnostic tests are utilized to identify the presence or absence of a condition in order to develop an appropriate treatment plan [12]. Many performance measures are used to evaluate a diagnostic test. These measures include sensitivity, specificity, false positive rate, false negative rate, Positive Predictive Value (PPV), Negative Predictive Value (NPV), positive likelihood ratio (LR+), negative likelihood ratio (LR-), accuracy, Youden Index (YI) and Diagnostic Odds Ratio (DOR) (Table 1 and Table 2) [13-15].

Meta-analysis

Meta-analysis is a statistical method that aims to provide more reliable and accurate findings via combining and summarizing the results from previous individual studies [3,16,17]. In 1954, Cochran developed a method for parameter estimation by bringing together researches made in different places, times and areas in an appropriate form [18]. Meta-analysis has been used in 1980s mostly to assess the clinical efficacy of individual medical interventions and since then, it has been a required and advocated statistical analysis in various disciplines [4].

In meta-analysis, different estimations are provided depending upon the contents of the study and these estimations are essential for determining the combined effect and assigning study weights. One of the models utilized in the meta-analysis is the fixed-effects model and the other is the random-effects model [9,16,17].

The fixed-effects model is based on the ground of the assumption that all studies included in the analysis predict the same effect size. In other words, it is assumed that if a trial has an effect, this effect does not interact with the study criteria and it remains constant. In the fixed-effects model, it is assumed that the differences between the effect sizes are the results of the sampling error. In this model, relatively narrower confidence intervals are obtained, accurate information about the homogeneity of the studies cannot be estimated since the between-study variance is not taken into account, and studies with small sample size may not be as sensitive as the ones with large samples [1,9,17,19,20].

The random-effects model makes calculations taking into consideration both the variances between the

Table 3: Estimation methods for meta-analysis.

Estimator	Abbreviation	Iterative/Non-iterative
Method of Moment Estimator		
DerSimonian and Laird	DL	Non-iterative
Hedges and Olkin	HO	Non-iterative
Paule and Mandel	PM	Iterative
Maximum Likelihood Estimators		
Restricted Maximum Likelihood	REML	Iterative
Maximum Likelihood	ML	Iterative
Model Error Variance Estimator		
Sidik and Jonkman	SJ	Non-iterative

studies and within each study. The random-effects model assumes that the heterogeneity of all effect sizes arises both from the sampling error and the variations within the study population. Since the between-study variances are taken into account with this model, the homogeneity of the studies can be assessed, and it is more sensitive in small sample sized studies [1,9,17,19,20].

Methods to estimate between-study variance

In the meta-analysis, there are various methods to estimate the between-study variance. Some of those are DerSimonian-Laird (DL), Restricted Maximum Likelihood (REML), Sidik and Jonkman (SJ), Hedges and Olkin (HO), Maximum Likelihood (ML), and Paule and Mandel (PM) estimation methods (Table 3) [3,9,10,21].

DerSimonian and Laird (DL) method: DL estimator is a non-iterative method that is frequently used as the default approach in many softwares [3,9]. τ^2 which is the between-study variance for random effect size model, and w_i which is the reverse of fixed effect variance for each study are used to calculate the new weights as

$$w_i^* = \frac{1}{[v_i + \hat{\tau}^2]}$$

From here $\hat{\tau}^2$,

$$\hat{\tau}^2 = \begin{cases} \frac{Q - (k-1)}{\sum_{i=1}^k w_i - \frac{\sum_{i=1}^k w_i^2}{\sum_{i=1}^k w_i}}, & Q \geq df \\ 0, & Q < df \end{cases}$$

When $\hat{\tau}^2$ equals to zero, it transforms from the random-effects model to the fixed-effects model.

$$Q = \sum_{i=1}^k w_i (\ln OR_i)^2 - \frac{(\sum_{i=1}^k w_i \ln OR_i)^2}{\sum_{i=1}^k w_i}$$

The above-mentioned Q value is calculated as

$$\ln \bar{T}_{DL} = \frac{\sum_{i=1}^k w_i^* \ln OR_i}{\sum_{i=1}^k w_i^*}$$

The combined effect size is calculated as

$$\ln \bar{T}_{DL} = \frac{\sum_{i=1}^k w_i^* \ln OR_i}{\sum_{i=1}^k w_i^*}$$

The variance of the combined estimation is calculated as

$$Var(\bar{T}_{DL}) = \frac{1}{\sum_{i=1}^k w_i^*}$$

and % $(1 - \alpha)$ the confidence interval is calculated as stated below

$$\exp \left[\ln(\bar{T}_{DL}) - z_{\frac{\alpha}{2}} \sqrt{Var(\bar{T}_{DL})} \right] \leq \theta \leq \exp \left[\ln(\bar{T}_{DL}) + z_{\frac{\alpha}{2}} \sqrt{Var(\bar{T}_{DL})} \right] \quad [17,22,23].$$

Restricted Maximum Likelihood (REML) method: REML estimation method is a well-known technique in the statistical literature and in this estimation method, the between-study variance (τ^2) is calculated via double-iterative

process. The first iteration includes the estimation of maximum likelihood estimator of $\hat{\tau}_{ML}^2$ [3,8,10]. The estimate of $\hat{\tau}_{REML}^2$ is obtained by the derivative of the restricted log-likelihood function.

$$\ln L(\tau^2) = -\frac{k}{2} \ln(2\pi) - \frac{1}{2} \sum_{i=1}^k \ln(v_i + \tau^2) - \frac{1}{2} \sum_{i=1}^k \frac{((y_i - \hat{\mu}_{RE}(\hat{\tau}_{ML}^2))^2)}{v_i + \tau^2} - \frac{1}{2} \ln \left(\sum_{i=1}^k \frac{1}{v_i + \tau^2} \right)$$

with respect to τ^2 equals to zero and the resulting solution of the equation for τ^2 is,

$$\hat{\tau}_{REML}^2 = \max \left\{ 0, \frac{\sum_{i=1}^k (w_{i,RE})^2 ((y_i - \hat{\mu}_{RE}(\hat{\tau}_{ML}^2))^2 - v_i)}{\sum_{i=1}^k (w_{i,RE})^2} + \frac{1}{\sum_{i=1}^k w_{i,RE}} \right\}$$

From here, it is provided via

$$w_{i,RE} = \frac{1}{v_i + \hat{\tau}_{REML}^2} \quad [9].$$

Sidik and Jonkman (SJ) method: This estimation method is proposed by Sidik and Jonkman and it is a non-iterative technique based on weighted least squares method [24]. To obtain the SJ estimator $\hat{\tau}_i = \frac{v_i}{\hat{\tau}_0^2}$ (assuming $\hat{\tau}_0^2 \neq 0$) and with this equation $\hat{q}_i = \hat{\tau}_i + 1$ values are calculated.

Here, $\hat{\tau}_0^2 = \frac{\sum_{i=1}^k (y_i - \bar{y})^2}{k}$ is the initial estimate of the between-study variance. Then, the SJ estimator is obtained by setting the quantity $\sum \hat{q}_i^{-1} (y_i - \hat{\mu}_{\hat{q},RE})^2$ equal to its expected value $\hat{\tau}_{SJ}^2 = \frac{1}{k-1} \sum_{i=1}^k \hat{q}_i^{-1} (y_i - \hat{\mu}_{\hat{q},RE})^2$ [9].

Hedges and Olkin (HO) method: Hedges and Olkin estimation method was first defined by Cochran [18]. Hedges (1983) discussed the estimation method for the between-study variance component in the meta-analytic context. The estimator is obtained by setting the sample variance

$$S_y^2 = \frac{1}{k-1} \sum_{i=1}^k (y_i - \bar{y})^2$$

equal to its expected value and solving τ^2 , which yields

$$\hat{\tau}_{HO}^2 = \max \left\{ 0, \frac{1}{k-1} \sum_{i=1}^k (y_i - \bar{y})^2 - \frac{1}{k} \sum_{i=1}^k v_i \right\} \quad [9].$$

Maximum Likelihood (ML) method: ML estimator is asymptotically efficient but it requires an iterative solution. According to the marginal distribution of $y_i \sim N(\mu, v_i + \tau^2)$, the estimation of $\hat{\tau}_{ML}^2$ is obtained by maximizing the log-likelihood function.

$$\ln L(\mu, \tau^2) = -\frac{k}{2} \ln(2\pi) - \frac{1}{2} \sum_{i=1}^k \ln(v_i + \tau^2) - \frac{1}{2} \sum_{i=1}^k \frac{(y_i - \mu)^2}{(v_i + \tau^2)}$$

Setting partial derivatives with respect to μ and τ^2 which are equal to zero, and solving the likelihood equations for the two parameters to be estimated, the ML estimators for μ and τ^2 can be obtained as follows

$$\hat{\mu}_{RE}(\hat{\tau}_{ML}^2) = \frac{\sum_{i=1}^k w_{i,RE} y_i}{\sum_{i=1}^k w_{i,RE}}$$

$$\hat{\tau}_{ML}^2 = \max \left\{ 0, \frac{\sum_{i=1}^k (w_{i,RE})^2 ((y_i - \hat{\mu}_{RE}(\hat{\tau}_{ML}^2))^2 - v_i)}{\sum_{i=1}^k (w_{i,RE})^2} \right\}$$

From here, it is provided via

$$w_{i,RE} = \frac{1}{v_i + \hat{\tau}_{ML}^2} \quad [9].$$

Paule and Mandel (PM) method: The Paule and Mandel estimation method has most of the advantages of the method of moments due to its' semiparametric characteristics and the lack of requirement of convergence diagnostics [7]. This method is essentially equivalent to the Empirical Bayes estimator discussed by Morris [9,25]. Using the random effect weights, this method is equivalent to empirical Bayes method. Paule and Mandel, proposed a special form of Q with a equation [26].

$$\hat{\tau}_{PM}^2 = \frac{Q_{gen} - \left(\frac{\sum_{i=1}^k a_i v_i - \frac{\sum_{i=1}^k a_i^2 v_i}{\sum_{i=1}^k a_i}}{\sum_{i=1}^k a_i - \frac{\sum_{i=1}^k a_i^2}{\sum_{i=1}^k a_i}} \right)}{\sum_{i=1}^k a_i - \frac{\sum_{i=1}^k a_i^2}{\sum_{i=1}^k a_i}}$$

$$a_i = w_{i,RE} = \frac{1}{v_i + \tau^2}.$$

The generalized Q statistic is

$$Q_{gen} = \sum w_{i,RE} (y_i - \hat{\mu}_{RE}(\tau^2))^2 \sim \chi_{k-1}^2 [9].$$

Implementation

In the application section of the study, we conducted a meta-analysis in order to evaluate the effectiveness of Clinical Oral Examination (COE) for predicting the diagnosis of oral dysplasia or Oral Squamous Cell Carcinoma (OSCC) of mucosal lesions that were submitted for biopsy and were diagnosed histologically. A Clinical Oral Examination (COE) is the principal strategy used to detect abnormal oral mucosal changes including OSCC and oral dysplasia, which is the initial stage of cellular transformation to malignancy [27]. Automated literature searches of articles published from Jan 1,

1966, through Jan 20, 2010, was completed by using the PubMed, Web of Knowledge and the Cochrane Library databases via using the search terms “oral mucosal lesion screening” and “oral lesions”. A total of 1,252 articles have met the inclusion criteria (1,195 studies in PubMed, 38 in the Cochrane Library and 19 in Web of Knowledge). Additional articles which included clinically detected lesions that were identified by means of visual examination and other visual techniques were also entered as subsets of data. In all enrolled studies, the main inclusion criterion was the presence of histological diagnoses which were obtained after tissue biopsy of clinically detected oral mucosal lesions.

In conclusion, twenty-four observational studies which included 7,079 patients and 1,956 biopsies met the inclusion criteria [27]. The analyses for diagnostic test and meta-analysis of the data were performed by using Open Meta-Analyst, R Packages, Meta Essential 1.4, STATA 13.0 statistical software.

Results and Discussion

First of all, the sensitivity, specificity, odds ratio, accuracy, Positive Likelihood Ratio (PLR), Negative Likelihood Ratio (NLR), Positive Predictive Value (PPV),

Table 4: Diagnostic test results of studies included in meta-analysis.

Author, Year	Sensitivity	Specificity	OR	Accuracy	PLR	NLR	PPV	NPV	Youden Index
Onofre, et al. (1997) [29]	0.938	0.808	63	0.83	4.875	0.077	0.5	0.984	0.745
Zheng, et al. (2002) [30]	0.988	0.016	1.328	0.569	1.004	0.756	0.57	0.5	0.004
Epstein, et al. (2003) [31]	0.981	0.033	1.759	0.634	1.015	0.577	0.638	0.5	0.014
Maraki, et al. (2004) [32]	0.789	0.139	0.607	0.265	0.917	1.512	0.181	0.733	-0.071
Ram & Siar (2005) [33]	0.98	0.062	3.267	0.758	1.045	0.32	0.766	0.5	0.043
Du, et al. (2007) [34]	0.985	0.984	4221	0.985	63.059	0.015	0.957	0.995	0.970
Epstein, et al. (2008) [35]	0.991	0.011	1.253	0.556	1.002	0.8	0.556	0.5	0.002
Allegra, et al. (2009) [36]	0.533	0.8	4.571	0.622	2.667	0.583	0.842	0.462	0.333
Wilder-Smith, et al. (2009) [37]	0.985	0.028	1.914	0.654	1.013	0.529	0.657	0.5	0.013
Arduino, et al. (2009) [38]	0.998	0.5	415	0.995	1.995	0.005	0.998	0.5	0.498
Nagaraju, et al. (2010) [39]	0.545	0.917	13.157	0.581	6.536	0.497	0.984	0.177	0.461
Koch, et al. (2011) [40]	0.996	0.007	1.711	0.63	1.003	0.586	0.631	0.5	0.003
Jerjes, et al. (2010) [41]	0.972	0.028	1	0.5	1	1	0.5	0.5	0
Prout, et al. (1997) [42]	0.5	0.524	1.1	0.522	1.05	0.955	0.091	0.917	0.024
Epstein H & N (2003) [43]	0.4	0.682	1.429	0.594	1.257	0.88	0.364	0.714	0.082
Remmerbach, et al. (2003) [44]	0.991	0.978	4815	0.987	45.574	0.009	0.991	0.978	0.969
Chen, et al. (2007) [45]	0.983	0.017	1	0.5	1	1	0.5	0.5	0
Bhalang, et al. (2008) [46]	0.986	0.025	1.872	0.649	1.012	0.541	0.652	0.5	0.011
Farah & Mccullo, et al. (2007) [47]	0.955	0.859	127.615	0.877	6.755	0.053	0.618	0.988	0.813
Mehrotra, et al. (2008) [48]	0.986	0.011	0.758	0.432	0.997	1.314	0.431	0.5	-0.003
McIntosh, et al. (2009) [49]	0.95	0.75	57	0.788	3.8	0.067	0.475	0.984	0.7
Mehrotra, et al. (2010) [50]	0.971	0.002	0.068	0.065	0.973	14.294	0.064	0.5	-0.027
Koch, et al. (2011) [51]	0.985	0.011	0.736	0.425	0.996	1.353	0.424	0.5	-0.004
Güneri, et al. (2011) [52]	0.964	0.468	23.727	0.622	1.812	0.076	0.45	0.967	0.432

Negative Predictive Value (NPV), and Youden Index (YI) were calculated (Table 4). When the Odds Ratio values (OR) of the studies were considered, both very high (OR = 4815) and very low OR values (OR = 0.068) were observed. The accuracy value, which is expected to be high in a favorable diagnostic test, has varied between 0.065 and 0.995 among the studies.

The Q test was utilized to evaluate the heterogeneity between studies. As a result, it was assessed that the studies were heterogeneous (Q = 68.94, p = 0.00 < 0.05). Thus, random effect model was used for meta-analysis.

Using Open Meta Analyst statistical software, the meta-analysis of the random effect model that was performed according to the DerSimonian and Laird (DL) estimation method revealed that the pooled sensitivity value of COE was high [0.953 (95% CI: 0.895-0.979)] and the pooled specificity was low [0.25 (95% CI: 0.124-0.44)] (Table 5).

When the PLR and NLR were considered, the pooled PLR value was 1.053 (95% CI: 1.00-1.11) and the pooled NLR was 0.469 (95% CI: 0.341-0.645). In general, a PLR value above 10.0 indicates that the test makes a significant contribution to the diagnostic process and a NLR below 0.2 indicates that the test is good at ruling out diseases [15,28]. Additionally, PLR and NLR values of 1 demonstrate that the test provides no information about the likelihood of the disease. In our study, the

pooled odds ratio (OR) was 6.031 (95% CI: 2.208-16.471), revealing the ineffectiveness of the COE in prediction of oral dysplasia or OSCC.

The results of the analyses obtained with DL, REML, ML, PM, HO, and SJ estimation methods using R, Open Meta Analyst, Meta Essential softwares are presented in Table 6. The DL estimation method was present in all the software programs used in the study. The Q statistic value that was calculated for evaluation of the homogeneity by using the DL method in R, Open Meta Analyst and Meta Essential softwares yielded to 68.943 (p < 0.0001), and the lowest I^2 and τ^2 values were obtained. Based on these results, it can be concluded that a moderate level of heterogeneity was present. In R and Open Meta Analyst softwares, Restricted Maximum Likelihood (REML), Maximum Likelihood (ML), Paule and Mandel (PM), Hedges and Olkin (HO) and Sidik and Jonkman (SJ) estimation methods were utilized and similar results were obtained with both softwares. According to the results, the highest I^2 and τ^2 values were obtained using the non-iterative SJ estimation method. The analysis with the PM estimation method which is simple and does not require distributional assumption, the lowest I^2 value was obtained following REML and ML estimation methods (I^2 = 72.80%).

The publication bias was investigated by using the Egger weighted regression method and a funnel plot chart was prepared (Table 7 and Figure 1). Egger regression method and the funnel plot chart showed that, with 95% confidence intervals, publication bias was not present (p = 0.087 > 0.05).

Conclusion

The results of our study indicate that Clinical Oral Examination (COE) is not a sufficient technique for the diagnosis. Except Der Simonian and Laird (DL)

Table 5: The results of meta-analysis.


Diagnostic Test Indicators	95% CI
Pooled Sensitivity: 0.953	(0.895-0.979)
Pooled Specificity: 0.25	(0.124-0.44)
Pooled Odds Ratio: 6.031	(2.208-16.471)
Pooled PLR +: 1.053	(1.00-1.11)
Pooled NLR-: 0.469	(0.341-0.645)

Table 6: Meta-analysis results of different estimation methods in different softwares.

Software	Estimation Methods	Pooled OR (95% CI)	Homogeneity (Q Test) p < 0.05	τ^2	I^2
R	DL	6.031 (2.208-16.471)	68.943 (p < 0.0001)	3.489	66.64%
	REML	6.098 (2.063-18.02)	68.943 (p < 0.0001)	4.461	71.86%
	ML	6.079 (2.112-17.5)	68.943 (p < 0.0001)	4.128	70.26%
	PM (EB)	6.108 (2.033-18.351)	68.943 (p < 0.0001)	4.676	72.80%
	HO	6.115 (2.011-18.593)	68.943 (p < 0.0001)	4.834	73.46%
	SJ	6.140 (1.915-19.682)	68.943 (p < 0.0001)	5.557	76.08%
Open Meta Analyst (OPENMEE)	DL	6.031 (2.208-16.471)	68.943 (p < 0.001)	3.489	66.63%
	REML	6.098 (2.063-18.02)	68.943 (p < 0.001)	4.461	71.86%
	ML	6.079 (2.112-17.5)	68.943 (p < 0.001)	4.128	70.26%
	PM (EB)	6.108 (2.033-18.352)	68.943 (p < 0.001)	4.676	72.80%
	HO	6.115 (2.011-18.593)	68.943 (p < 0.001)	4.834	73.45%
	SJ	6.140 (1.915-19.682)	68.943 (p < 0.001)	5.557	76.08%
Meta Essential	DL	6.03 (1.91-19.09)	68.94 (p < 0.000)	3.49	66.64%

Table 7: Egger regression test.

	Coef.	Std. Error	t	p	95% Confidence Interval	
					LL	UL
Slope	-0.105	0.828	-0.13	0.9	-1.824	1.613
Bias	1.187	0.663	1.79	0.087	-0.188	2.563



estimation methods, analyses could be performed with other estimation methods in ready-made softwares. Furthermore, it can be concluded that the appropriate software program for meta-analysis varies depending on the user's needs and preferences.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Kurt S (2009) Meta Analizin Klinik Çalışmalarda Kullanımı Üzerine Bir Uygulama. Yüksek Lisans Tezi, Marmara Üniversitesi Ekonometri Anabilim Dalı, İstanbul, 77.
- Karahan AE (2014) Hayvancılıkta Meta Analiz Uygulamaları. Yüksek Lisans Tezi, Ege Üniversitesi Zootekni Anabilim Dalı, İzmir, 53s.
- Langan D, Higgins JPT, Jackson D, Bowden J, Veroniki AA, et al. (2018) A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses. *Res Synth Methods* 10: 83-98.
- Crombie IK, Davies HT (2009) What is meta-analysis? Hayward Medical Communications.
- Çubuk M (2011) Tani Doğruluğunu Ölçülerini Etkileyen Faktörlerin Belirlenmesi. Hacettepe Üniversitesi Biyoistatistik Anabilim Dalı, Ankara, 107.
- Viechtbauer W, Lopez-Lopez JA, Sanchez-Meca J, Marin-Martinez F (2014) A comparison of procedures to test for moderators in mixed-effects meta-regression models. *Psychol Methods* 20: 360-374.
- Jackson D, Veroniki AA, Law M, Tricco AC, Baker R (2017) Paule-Mandel estimators for network meta-analysis with random inconsistency effects. *Res Synth Methods* 8: 416-434.
- Langan D, Higgins JPT, Simmonds M (2017) Comparative performance of heterogeneity variance estimators in meta-analysis: A review of simulation studies. *Res Synth Methods* 8: 181-198.
- Veroniki AA, Jackson D, Viechtbauer W, Bender R, Bowden J, et al. (2016) Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Methods* 7: 55-79.
- Thorlund K, Wetterslev J, Awad T, Thabane L, Gluud C (2011) Comparison of statistical inferences from the DerSimonian-Laird and alternative random-effects model meta-analyses - an empirical assessment of 920 Cochrane primary outcome meta-analyses. *Res Synth Methods* 2: 238-253.
- Bowden J, Tierney JF, Copas AJ, Burdett S (2011) Quantifying, displaying and accounting for heterogeneity in the meta-analysis of RCTs using standard and generalised Q statistics. *BMC Med Res Methodol* 11: 41.
- White S, Schultz T, Enuneh YAK (2011) Synthesizing evi-

- dence of diagnostic accuracy. Lippincott Williams &Wilkins, JBI Book Collection, Philadelphia, 116.
13. Aslan D, Sandberg S (2007) Simple statistics in diagnostic tests. *JMB* 26: 309-313.
 14. Glas AS, Lijmer JG, Prins MH, Bossel GJ, Bossuyt PMM (2003) The diagnostic odds ratio: A single indicator of test performance. *J Clin Epidemiol* 56: 1129-1135.
 15. Karaağoğlu E, Karakaya J, ve Kiliçkap M (2016) Tani Testlerinin Değerlendirilmesinde İstatistiksel Yöntemler. Ankara, 261.
 16. Borenstein M, Hedges LV, Higgins J, Rothstein H (2009) Introduction to meta-analysis. John Wiley & Sons, Ltd, Chichester, UK.
 17. Yeniay H (2013) Yaşam Verilerinin Meta Analizi. Yüksek Lisans Tezi, Hacettepe Üniversitesi, İstatistik Anabilim Dalı, Ankara, 72.
 18. Cochran WG (1954) The combination of estimates from different experiments. *Biometrics* 10: 101.
 19. Kinay E (2012) Üniversite Giriş Sınavı Yordama Yeterliliği Çalışmalarının Meta Analizi. Yüksek Lisans Tezi, Ankara Üniversitesi Eğitim Bilimleri Enstitüsü, Ankara, 110.
 20. Küçükönder H (2007) Meta Analiz ve Tarımsal Uygulamalar, Yüksek Lisans Tezi, Kahramanmaraş Sütçü İmama Üniversitesi Zootekni Anabilim Dalı, Kahramanmaraş, 100.
 21. Panityakul T, Bumrungrsup C, Knapp G (2013) On estimating residual heterogeneity in random-effects meta-regression: A comparative study. *J Stat Theory Appl* 12: 253-265.
 22. DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Control Clin Trials* 7: 177-188.
 23. Petitti DB (2000) Meta-Analysis, Decision Analysis, and Cost-Effectiveness Analysis. Oxford University Press, New York, 306.
 24. Sidik K, Jonkman JN (2005) Simple heterogeneity variance estimation for meta-analysis. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 54: 367-384.
 25. Morris CN (1983) Parametric empirical Bayes inference: Theory and applications. *Journal of the American Statistical Association* 78: 47.
 26. Paule RC, Mandel J (1982) Consensus values and weighting factors. *Journal of Research of the National Bureau of Standards* 87: 377-385.
 27. Epstein JB, Güneri P, Boyacıoğlu H, Abt E (2012) The limitations of the clinical oral examination in detecting dysplastic oral lesions and oral squamous cell carcinoma. *J Am Dent Assoc* 143: 1332-1342.
 28. Deeks JJ (2001) Systematic reviews in healthcare: Systematic reviews of evaluations of diagnostic and screening tests. *BMJ* 323: 157-162.
 29. Onofre MA, Sposto MR, Navarro CM, Motta ME, Turatti E, et al. (1997) Potentially malignant epithelial oral lesions: Discrepancies between clinical and histological diagnosis. *Oral Dis* 3: 148-152.
 30. Zheng W, Soo KC, Sivanandan R, Olivo M (2002) Detection of squamous cell carcinomas and pre-cancerous lesions in the oral cavity by quantification of 5-aminolevulinic acid induced fluorescence endoscopic images. *Lasers Surg Med* 31: 151-157.
 31. Epstein JB, Zhang L, Poh C, Nakamura H, Berean, K, et al. (2003) Increased allelic loss in toluidine blue-positive oral premalignant lesions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 95: 45-50.
 32. Maraki D, Becker J, Boecking A (2004) Cytologic and DNA-cytometric very early diagnosis of oral cancer. *J Oral Pathol Med* 33: 398-404.
 33. Ram S, Siar CH (2005) Chemiluminescence as a diagnostic aid in the detection of oral cancer and potentially malignant epithelial lesions. *Int J Oral Maxillofac Surg* 34: 521-527.
 34. Du GF, Li CZ, Chen HZ, Chen XM, Xiao Q, et al. (2007) Rose bengal staining in detection of oral precancerous and malignant lesions with colorimetric evaluation: A pilot study. *Int J Cancer* 120: 1958-1963.
 35. Epstein JB, Silverman S Jr, Epstein JD, Lonky SA, Bride MA (2008) Analysis of oral lesion biopsies identified and evaluated by visual examination, chemiluminescence and toluidine blue. *Oral Oncol* 44: 538-544.
 36. Allegra E, Lombardo N, Puzzo L, Garozzo A (2009) The usefulness of toluidine staining as a diagnostic tool for precancerous and cancerous oropharyngeal and oral cavity lesions. *Acta Otorhinolaryngol Ital* 29: 187-190.
 37. Wilder-Smith P, Lee K, Guo S, Zhang J, Osann K, et al. (2009) In vivo diagnosis of oral dysplasia and malignancy using optical coherence tomography: Preliminary studies in 50 patients. *Lasers Surg Med* 41: 353-357.
 38. Arduino PG, Surace A, Carbone M, Elia A, Massolini G, et al. (2009) Outcome of oral dysplasia: A retrospective hospital-based study of 207 patients with a long follow-up. *J Oral Pathol Med* 38: 540-544.
 39. Nagaraju K, Prasad S, Ashok L (2010) Diagnostic efficiency of toluidine blue with Lugol's iodine in oral premalignant and malignant lesions. *Indian J Dent Res* 21: 218-223.
 40. Koch FP, Kunkel M, Biesterfeld S, Wagner W (2011) Diagnostic efficiency of differentiating small cancerous and precancerous lesions using mucosal brush smears of the oral cavity - A prospective and blinded study. *Clin Oral Investig* 15: 763-769.
 41. Jerjes W, Upile T, Conn B, Hamdoon Z, Betz CS, et al. (2010) In vitro examination of suspicious oral lesions using optical coherence tomography. *Br J Oral Maxillofac Surg* 48: 18-25.
 42. Prout MN, Sidari JN, Witzburg RA, Grillone GA, Vaughan CW (1997) Head and neck cancer screening among 4611 tobacco users older than forty years. *Otolaryngol Head Neck Surg* 116: 201-208.
 43. Epstein JB, Feldman R, Dolor RJ, Porter SR (2003) The utility of toluidine chloride rinse in the diagnosis of recurrent or second primary cancers in patients with prior upper aerodigestive tract cancer. *Head Neck* 25: 911-921.
 44. Remmerbach TW, Weidenbach H, Müller C, Hemprich A, Pomjanski N, et al. (2003) Diagnostic value of nucleolar organizer regions (AgNORs) in brush biopsies of suspicious lesions of the oral cavity. *Anal Cell Pathol* 25: 139-146.
 45. Chen Y-W, Lin J-S, Fong JH-J, Wang I-K, Chou S-J, et al. (2007) Use of methylene blue as a diagnostic aid in early detection of oral cancer and precancerous lesions. *Br J Oral Maxillofac Surg* 45: 590-591.
 46. Bhalang K, Suesuwan A, Dhanuthai K, Sannikorn P, Luangjarmekorn L, et al. (2008) The application of acetic acid in the detection of oral squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 106: 371-376.
 47. Farah CS, McCullough MJ (2007) A pilot case control study on the efficacy of acetic acid wash and chemiluminescent illumination (ViziLite) in the visualisation of oral mucosal white lesions. *Oral Oncol* 43: 820-824.

48. Mehrotra R, Singh MK, Pandya S, Singh M (2008) The use of an oral brush biopsy without computer-assisted analysis in the evaluation of oral lesions: A study of 94 patients. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 106: 246-253.
49. McIntosh L, McCullough MJ, Farah CS (2009) The assessment of diffused light illumination and acetic acid rinse (MicroLux/DL) in the visualisation of oral mucosal lesions. *Oral Oncol* 45: e227-e231.
50. Mehrotra R, Singh M, Thomas S, Nair P, Pandya S, et al. (2010) A cross-sectional study evaluating chemiluminescence and autofluorescence in the detection of clinically innocuous precancerous and cancerous oral lesions. *J Am Dent Assoc* 141: 151-156.
51. Koch FP, Kaemmerer PW, Biesterfeld S, Kunkel M, Wagner W (2011) Effectiveness of autofluorescence to identify suspicious oral lesions - A prospective, blinded clinical trial. *Clin Oral Investig* 15: 975-982.
52. Güneri P, Epstein JB, Kaya A, Veral A, Kazandi A, et al. (2011) The utility of toluidine blue staining and brush cytology as adjuncts in clinical examination of suspicious oral mucosal lesions. *Int J Oral Maxillofac Surg* 40: 155-161.