



REVIEW ARTICLE

Effectiveness of Bright Light Therapy in Managing Behavioral and Psychological Symptoms of Dementia for Patients with Dementia: An Integrative Literature Review

Chan Shi Qing Charlotte* and Tay Swee Cheng Rosy

Woodlands Health, University of Glasgow, Singapore



*Corresponding author: Chan Shi Qing Charlotte, Woodlands Health, Singapore, Tel: +65-88762108

Abstract

Background: Dementia presents a significant public health concern, with a high prevalence rate and considerable challenges in managing behavioural and psychological symptoms of dementia (BPSD) for patients with dementia (PwD).

Aim: The aim of this integrative literature review was to evaluate the effectiveness of BLT in managing BPSD for PwD.

Methods: A literature search was conducted using PubMed, CINAHL, and Web of Science. Studies were included if they involved people with dementia of any type receiving BLT of any methods, forms, duration, intensity and frequency, and had outcomes measures including BPSDs such as sleep disturbance, cognition, depression, and/or agitation. The Joanna Briggs Institute critical appraisal checklists were used to evaluate the methodological quality of the studies. Seven studies were included in the final review.

Results: Three themes were identified: The impact of BLT on sleep, the impact of BLT on mood and behaviour, and the impact of BLT on cognition. Studies found varied results, with subjective sleep showing consistent improvements but objective measures having mixed findings. BLT showed potential in enhancing mood and behaviour, especially in advanced dementia stages, though outcomes depended on dementia type and severity. Notably, BLT consistently improved cognitive function across studies, suggesting promise as a non-pharmacological intervention for cognitive enhancement in PwD.

Conclusion: The use of BLT in managing BPSD in PwD appears promising. Nevertheless, further research may be necessary to optimise the effectiveness of BLT across varied disease stages and types and strengthen the evidence for BLT as an effective intervention.

Keywords

Light therapy, BLT, BPSD, PwD, Dementia, Nursing, Behavioural, Psychological, Cognition

Introduction

The World Health Organization (WHO) [1] has reported a global increase in the proportion of older adults within populations across countries worldwide, where one in six people globally will be above the age of 60 years by 2030. As the aging population continues to increase, individuals are at a higher risk of experiencing complex health conditions often referred to as geriatric syndromes, such as dementia, depression, diabetes, hearing loss, cataracts, and osteoarthritis [1].

Dementia, a neurocognitive brain disorder, presents with cognitive and non-cognitive symptoms impacting daily functioning [2]. According to WHO [3], dementia is a major health problem among the elderly population globally, leading to disability, dependence, and mortality. With the aging population worldwide, there is an expected rise in the incidence of dementia, projected to increase from an estimated 57 million to 153 million cases by 2050 [3]. While age remains the primary risk factor of dementia, disparities in prevalence linked to gender are predominantly attributed to the longer life expectancy of females [4]. Furthermore, during their illness trajectory, approximately 80% of PwD will develop BPSD, with 97% developing at least one symptom within five years [5]. BPSD includes disruptive

behaviour, mood, cognition, or perception symptoms such as agitation, irritability, anxiety, depression, apathy, and sleep disturbances [6].

In the 1940s, symptoms now defined as BPSD were observed to worsen during the late afternoon or early evening [7]. Thus, it was theorised that the primary reason for this phenomenon was the difficulty in retaining a spatial image under reduced light conditions [7]. This relation was subsequently termed “sundowning”, attributing the behaviour changes to the decreased light levels when the sun sets [8]. While the concept of sundowning syndrome is commonly used today, its definition remains unclear, and the underlying mechanism is not fully understood [8]. Evidence suggested that the underlying reason could be due to the dysregulation of circadian biology and alternations in circadian rhythms associated with the decreased light levels [9]. Furthermore, according to current clinical and preclinical research, the circadian rhythm, which is influenced by light levels may have an impact on mood, sleep, cognition, and aggressive behaviour [10]. Consequently, inappropriate light levels and wrestling dysregulated circadian rhythms could potentially be a plausible explanation for the sundowning syndrome [10].

BPSD can interfere with clinical management, causing distress to patients and caregivers, and is associated with diminished quality of life [11]. PwD has a higher risk of institutionalisation and the presence of BPSD can further complicate the treatment due to the special treatment requirements and additional behavioural and medical complications, leading to extended hospitalisation stays and increased costs [12]. In clinical settings, BPSD inflicts greater stress on caregivers and is commonly described as more challenging to manage than the cognitive deficits typically associated with dementia with agitation and sleep disturbances as prominent symptoms [12]. BPSD symptoms pose complications during routine nursing tasks including issues such as catheter removal and food throwing, and can lead to physical and verbal incidents for nursing staff during the delivery of nursing care, contributing to job dissatisfaction, burnout, and increased attrition rates among nursing staff [12-15].

Current BPSD treatment options are limited, raising concerns about side effects and insufficient research support [15]. In clinical settings, antipsychotic medication are commonly used for the management of BPSD but have limited effectiveness and significant adverse side effects such as the increased risk of hospitalisation, falls, and mortality [16]. Therefore, guidelines from experts favour non-pharmacological interventions for BPSD due to their significant influence on overall BPSD outcomes and the absence of adverse events [17]. However, restraints, one of the most commonly used non-pharmacological interventions to

manage BPSD, have been proven to further exacerbate BPSD and lead to numerous physical injuries [18]. Furthermore, healthcare providers often face emotional conflict regarding the use of restraints and their potential dehumanising impact on the patient’s dignity and autonomy [19]. Therefore, non-pharmacological treatment involving environmental and behavioural interventions such as bright light therapy could potentially serve as first-line alternatives.

BLT offers a feasible non-pharmacological treatment for BPSD in PwD, yielding positive outcomes in sleep, agitation, and cognitive functioning [20]. Older adults with dementia experience reduced general sensory input, demonstrating decreased sensitivity to the impact of light on the suprachiasmatic nucleus (SCN), and limited exposure to bright environmental light [21]. BLT can potentially reverse degenerative alterations in the SCN, which form the biological foundation for circadian rhythm disturbances such as sleep-wake rhythm and rest-activity in PwD [21]. Most clinical settings maintain consistently low light levels which might not provide an adequate stimulus for the circadian clock to synchronise with the 24-hour day [21]. The appropriate light dose can effectively influence circadian rhythms, thereby improving factors such as sleep quality, depression, functional status, or behavioural issues in PwD [20]. BLT involves controlled light delivered through various means such as natural outdoor sunlight, or artificial indoor light sources such as light boxes, ceiling lights, or stimulation dawn-dusk illumination [22]. Furthermore, it can be administered at different times of the day such as in the morning, afternoon, or intermittently throughout the day, to prevent the patients from sleeping during those periods [23]. Moreover, BLT is non-invasive and has minimal side effects, where users can engage in various activities within their field of vision while undergoing BLT, making it an ideal non-pharmacological intervention easily implementable across institutional settings [24].

Since the 1900s, BLT has been a subject of research in managing BPSD in PwD, particularly in sleep and circadian rhythms [25]. While some reviews have investigated its effects on cognition, mood, and behaviour, the reported findings are inconsistent. The identification of an effective and feasible non-pharmacological treatment, such as BLT, for managing BPSD in PwD is imperative for tailored nursing care. Utilising such behaviour management intervention can enhance the individual’s quality of life, and ensure patient safety during the nursing care delivery. Additionally, introducing BLT can potentially reduce the stress and burden associated with caregivers for informal caregivers such as family members and formal caregivers such as nurses. Nursing management can consider instituting BLT as part of policy changes in improving holistic care of PwD and nurses’ job satisfaction. Therefore, this review seeks to offer current insights and robust evidence regarding

Table 1: Inclusion and exclusion criteria.

	Inclusion Criteria	Exclusion Criteria
Population	<ul style="list-style-type: none"> - Participants with any form of dementia - Participants currently residing in clinical settings 	<ul style="list-style-type: none"> - Participants with other neurologic conditions - Participants in home settings
Intervention	<ul style="list-style-type: none"> - BLT of any methods, forms, duration, intensity and frequency 	<ul style="list-style-type: none"> - Other non-pharmacological interventions such as music therapy, art therapy
Outcome	BPSD	Symptoms non-related to BPSD
Publication type	Primary research articles: <ul style="list-style-type: none"> - Qualitative: RCTS, Quasi-experimental - Quantitative 	Secondary Sources
Language	English language	Non-English language
Publication Period	Published between 2019-2023	Published before 2019
Text Availability	Full-text articles	Non-full text articles

the effectiveness of BLT in managing BPSD in PwD, facilitating the development of new knowledge for informed clinical practice, enhancing the quality of life in PwD, and improving dementia care.

Methods

The literature search was done following the PRISMA 2020 guidelines [26]. Three databases (PubMed, CINAHL, and Web of Science) were searched using keywords such as: “dementia”, “light therapy”, “behavioural”, “psychological”, and “cognition” with the use of Boolean operators to identify relevant studies. Search filters were applied across the databases to obtain studies that were published between 2019-2023, and in English.

A total of 81 articles were identified from the databases. Mendeley Reference Manager was used for the removal of 12 duplicates gathered from the search results [27].

The remaining articles were subjected to title and abstract screening. Studies were included if they involved people with dementia of any type receiving BLT of any methods, forms, duration, intensity and frequency, and had outcomes measures including BPSDs such as sleep disturbance, cognition, depression, and/or agitation. Research articles that did not match the aims of the present literature review, articles focusing on participants with other neurological conditions, and studies that included other non-pharmacological interventions were excluded, together with non-full text and non-English articles published before 2019 (Table 1). After screening, 11 articles remained for full-text review. Two articles were excluded as one did not qualify as a primary studies while the other was conducted in a home setting. Ultimately, nine articles were considered eligible for this review [20,25,28-34]. Figure 1 shows a PRISMA flowchart illustrating the literature search flow.

Data Evaluation

The nine studies were evaluated for their

methodological quality using the Joanna Briggs Institute (JBI) critical appraisal tools. According to the JBI [35] level of evidence for effectiveness, all nine RCTs in this review were classified as Level 1.c, denoting the highest quality of evidence. While JBI levels of evidence offer valuable ranking, it is not a definitive measure of study quality and should not preclude the need for critical appraisal and clinical reasoning when applying evidence [36]. Therefore, the methodological quality of the nine studies will be further evaluated using the JBI checklist for randomised clinical trials [37], critically appraising the sample size, population characteristics, validity and reliability of measurement tools, data collection method, study design, and risk of bias. Each study was given a quality percentage by first scoring it out of 13 before converting the score into a percentage. Scores were obtained by summing the number of items on the checklist with a “Yes” response. The quality appraisal are summarised in Table 2.

Notably, all nine studies had a relatively small sample size of less than 50 participants and power analysis was implemented in only four studies [20,25,30,32], with all recruiting fewer participants than calculated. Consequently, these small sample sizes could potentially compromise the robustness and generalisability of the results, resulting in insufficient statistical power to accurately detect the true effects of BLT. Additionally, a gender bias was evident, with a higher proportion of females, which could confound the result due to gender-specific responses to the light interventions [38]. Furthermore, while concerns were raised regarding the validity of watch actigraphy for objective sleep measurement, the homogeneity and reliability of the remaining measurement tools, assessing subjective sleep, mood and behaviour, and cognition, were deemed valid and reliable. Lastly, despite the potential risk of biases in majority of the studies, the lack of allocation concealment and participant blinding is unlikely to undermine the validity




Table 2: Joanna Briggs Institute critical appraisal checklist for randomised clinical trials.

Article	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13	Score
Cremascoli, et al. [30]	Y	Y	Y	N	Y	Y	Y	Y	N	Y	N	N	Y	9/13 (69.2%)
Figueiro, et al. [31]	Y	N	U	N	Y	Y	Y	Y	N	Y	N	N	Y	7/13 (53.8%)
Kim, et al. [34]	Y	N	Y	U	U	Y	U	Y	N	Y	N	Y	Y	7/13 (53.8%)
Cibeira, et al. [29]	U	N	N	N	N	N	N	N	N	N	N	Y	N	1/13 (7.69%)
Hjetland, et al. [32]	Y	N	Y	N	Y	Y	Y	Y	N	Y	N	Y	Y	9/13 (69.2%)
Kolberg, et al. [25]	Y	N	Y	N	Y	Y	Y	Y	N	Y	N	Y	Y	10/13 (76.9%)
Kolberg, et al. [20]	Y	N	Y	N	Y	Y	Y	Y	Y	Y	N	Y	Y	9/13 (69.2%)
Bromundt, et al. [28]	N	N	Y	N	N	N	N	Y	N	Y	N	N	N	3/13 (23%)
Kheradmand, et al. [33]	Y	Y	Y	U	U	Y	U	Y	N	Y	U	N	Y	7/13 (53.8%)

Note: Q1: Randomization; Q2: Concealed allocation; Q3: Groups similar at baseline; Q4: Participants blinded; Q5: Allocator blinded; Q6: Treatment Groups treated equally; Q7: Assessors blinded; Q8: Outcomes measured in the same way in groups; Q9: Outcomes measured reliably; Q10: Follow-up complete; Q11: Participants analysed in original group; Q12: Appropriate statistical analysis; Q13: Appropriate trial design. Abbreviations: Y: Yes; N: No; U: Unclear; N/A: not applicable

of the results or introduce significant bias due to the nature of the light intervention. However, two studies failed to demonstrate true RCT due to the absence of proper randomisation, lacking an appropriate washout period in their crossover RCT [28], and conducting a single-arm intervention study without a control group [29]. These deviations from the standard RCTs were not accounted for in the conduct and analysis which, therefore, introduced high risks of biases. This, in turn, suggests that the reliability and validity of the findings of both studies could be compromised and the evidence may not accurately reflect the true effects of the study, overall indicating a low quality of evidence. Consequently, these two studies were eventually excluded due to poor methodological quality [28,29].

Overall, based on the assessment of methodological rigor, four of the seven studies achieved a score of 70% and above, indicating high-quality evidence [20,25,30,32]. Among the remaining five studies, three attained at least 50% of the predefined appraisal criteria [31,33,34], indicating moderate-quality evidence. The two studies that were excluded, scored below 25%, indicating low-quality evidence [28,29].

Results

Characteristics of studies

The main characteristics of the studies are summarised in Table 3. All seven studies in this review were RCTs, including the standard RCTs [30,33,34],

Table 3: Characteristics of studies.

Author/ Year	Country	Study Design, Setting, Period	Population (Sample)	Intervention Characteristics	Outcome Measured	Outcomes and Findings
Cremascoli, et al. [30]	Italy	Randomised placebo- controlled, single-blind trial, over a 3-month period at the Unit of Sleep Medicine of IRCCS Mondino Foundation	Patients with mild/moderate AD Total sample: n = 13 Intervention group: n = 8 Control group: n = 5 Mean age: 73 4 females, 9 males	Intervention group: Blue enriched light (Luminette glasses), 10,000 lux, 20 min once a day, every day, 4 weeks Control Group: Use of the same Luminette device set at a fluorescence of 50 lux, 20 min once a day, everyday, 4 weeks Light delivery was planned according to the individual circadian phase of each participants.	Sleep-wake rhythms, sleep quality, mood, and cognition Subjective sleep measurement: PSQI, ESS Objective sleep measurement: Actigraphy watch, 24h wake-sleep diaries Cognition: MMSE Mood: NPI, Beck depression inventory	The treatment induced an objective circadian phase shift consistent with the melatonin phase response curve to light exposure, led to a shortening of the phase angle DLMO-falling asleep time, and was associated with an improvement in subjective sleep quality and cognitive performance.
Figueiro, et al. [31]	USA	Randomised placebo- controlled, crossover design clinical trial 8 long-term care facilities for two 4- week period (separated by a 4-week washout)	Patients with moderate to late-stage AD and related dementia (ADRD) Total sample: n = 46 Intervention group: n = 25 Control group: n = 21 Mean age: 85.1 30 females, 16 males	Intervention group: High circadian stimulus, Full spectrum white light, 550-700 lux Bright light (light box, light panels) 24 h (tailored lighting), everyday, 14 weeks Control group: Low circadian stimulus, 110-200 lux dim light, 24h, everyday, 14 weeks Timers controlled all lights for both interventions, activating the lights according to the individual participants habitual wake times	Sleep quality, depression, agitation, quality of life Subjective sleep measurement: PSQI Objective sleep measurement: Actigraphy Mood and behaviour: CMAI, CSSD	The active intervention significantly improved Pittsburgh Sleep Quality Index scores compared to the active baseline and control intervention. The active intervention also resulted in significantly greater active versus control differences in intraday variability. As for secondary outcomes, the active intervention resulted in significant improvements in Cornell Scale for Depression in Dementia scores and significantly greater active versus control differences in Cohen-Mansfield Agitation Inventory scores

Hjetland, et al. [32]	Norway	24-week cluster randomised Placebo- controlled trial	<p>Patients with severe dementia Total sample: n = 69 Intervention Group: n = 33 Control group: n = 36 Mean age: 84 years females, 22 males</p>	<p>Intervention Group: ambient light of 1000 lx and 6000 K from 10:00 to 15:00, with gradually increasing and decreasing light levels prior to and following this interval, using ceiling mounted light-fixtures and light emitting diode technology</p> <p>Control Group: The placebo condition had continuous standard light levels (150-300 lx, ~ 3000 K)</p>	<p>Sleep outcomes</p> <p>Subjective Sleep: SDI</p> <p>Objective Sleep: actigraphy</p>	<p>Sleep as measured by the SDI was significantly improved in the intervention group compared to the control group from baseline to week 16 and from baseline to week 24. There was no significant effects in terms of actigraphically measured sleep.</p> <p>There was no effect according to the SDI at week 8 compared to the control group, the intervention group fared better from baseline to week 16 and from baseline to week 24, as measured by the proxy-rated SDI. Overall, proxy-rated sleep improved among nursing home patients with dementia following 16 and 24 weeks of BLT.</p> <p>These improvements were not corroborated by actigraphy recordings</p>
Kheradmand, et al. [33]	Iran	Double-blinded randomized placebo- controlled study	<p>Patients with moderate to severe AD Total sample: n = 32 Intervention group: 16 Control group: 16 Mean age: 77.13 females, 15 men</p>	<p>Intervention group: LLT for 10 minutes and 6 sessions (3 times a week for 2 weeks). The LLT system includes a helmet and a control device. Lasers operated in pulse mode with a frequency of 75 Hz. In one session, the cumulative dose for each laser emitter and all 24 lasers inside the helmet were 12 and 288 J</p> <p>Control group: Sham therapy for 10 minutes and 6 sessions (switched off laser)</p>	<p>Cognition: MMSE</p>	<p>The results showed considerable progress in the cognitive function of the patients undergoing LLT based on the MMSE results 2 weeks after initiation of treatment. In contrast, the MMSE scores worsened in the sham group. In addition, after 6 weeks, the positive effect of LLT on the cognitive function of the patients was also observed.</p>

Kim, et al. [34]	South Korea	Single-blind RCT	<p>Patients with mild and moderate AD Total sample: n = 25 Intervention group: n = 14 (77.36 ± 5.79 years; M: F = 2:12) Control group: n = 11 (78.55 ± 7.71 years; M:F = 5:6)</p>	<p>Intervention Group: Blue-enriched white light: 30 lux; Blue attenuating light (10 lx), 1 times/day, 7 d/week, 60 min, 2 weeks Control group: Same, blue-enriched white light + blue-attenuating sunglasses 1 times/day, 7 d/week, 60 min, 2 weeks.</p> <p>Both groups Sat about 60 cm away from a small LED light box for 1h each morning for 2 weeks.</p>	<p>Sleep, cognition, depression, mood, and behavior</p> <p>Sleep: PSQI, actigraphy, sleep diary</p> <p>Mood and behaviour: KNPI-Qs, CSDD-K, VAS</p> <p>Cognition: MMSE- KC</p>	<p>Blue-enriched LT had a significantly better effect on the Pittsburgh Sleep Quality Index at T2 compared to blue-attenuated LT, and a trend of better effectiveness on total sleep time at T2.</p> <p>There was a significant increase in Mini-Mental State Examination score at T2 after blue-enriched LT than that at T0.</p> <p>Our findings suggest that morning blue-enriched LT has a benefit in improving sleep and cognitive function in AD patients.</p>
Kolberg, et al. [25]	Bergen, Norway	24-week cluster randomised controlled trial	<p>Patients with severe dementia Total Sample: n = 61 Intervention Group = 31 Control group = 30 Mean age: 84 44 females, 20 males</p>	<p>Intervention Group: ceiling-mounted LED panels providing varying illuminance and correlated color temperature throughout the day, with a peak of 1000 lx, 6000 K between 10 a.m. and 3 p.m</p> <p>Control Group: lights bulbs were changed but still delivered standard indoor illumination (~150-300 lx, 3000 at eye level in the center of the room).</p>	<p>Rest-activity rhythms measured by actigraphs</p>	<p>After 16 weeks, the acrophase was advanced from baseline in the intervention group compared to the control group. There was no significant difference between the groups on any other rest-activity measures.</p> <p>When comparing parametric and non-parametric indicators of rest-activity rhythms, 25 out of 35 comparisons were significantly correlated. The present results indicate that ambient bright light treatment did not improve rest-activity rhythms for people with dementia.</p>

Table 4: Thematic analysis.

Author	Theme 1: Impact of BLT on sleep	Theme 2: Impact of BLT on Mood and Behaviour	Theme 2: Impact of BLT on Cognition
Cremaşcoli, et al. [30]	✓	✓	✓
Figueiro, et al. [31]	✓	✓	X
Hjetland, et al. [32]	✓	X	X
Kheradmand, et al. [33]	X	X	✓
Kim, et al. [34]	✓	✓	✓
Kolberg, et al. [25]	✓	X	X
Kolberg, et al. [25]	X	✓	X

cluster RCTs [20,25,32] and crossover RCTs [31]. Six studies were conducted in Western countries including Italy [30], Norway [20,25,32], the United States of America [31], and Iran [33], while one was conducted in Asia, specifically South Korea [34]. A total of 315 participants were included, with the largest group consisting of 69 participants [25,32]. The majority of participants were females, and the mean age of the participants was 81 years. Four studies only included patients with AD [30,31,33,34], while the remaining included participants with several types of dementia such as VD, Lewy Body Dementia and Frontotemporal Dementia [20,25,32]. Most studies included individuals with moderate to severe dementia [20,25,31-33].

The intervention characteristics varied in the type of light, intensity, duration, and frequency. Among the seven studies, BLT interventions can be grouped into three main categories: standard visible light, light-emitting diodes (LED) light, and laser light. Most of the studies employed standard visible light interventions, such as normal bright light [31,32] and blue-enriched light [30,34]. In contrast, three studies incorporated LED light sources, including LED helmet devices and ceiling-mounted LED panels [20,25], while the remaining study used low-level laser therapy [33]. BLT exposure duration ranged from ten minutes to 24 hours, spanning from three to seven days a week, and lasting two to 24 weeks. The intensity of BLT ranged from 550 lux to 10,000 lux. In three studies, the delivery of light intervention and illuminance was tailored according to each participant's circadian phase [30,31], while in another three studies, the light intervention provided varying illuminance throughout the day [20,25,32]. The remaining study involved participants receiving the light intervention at the same intensity for only one hour each morning [34]. Control conditions generally implemented low light intensities, ranging from 30 lux to 300 lux, with some studies requiring participants to wear blue-attenuating sunglasses [34] or turning the light intervention off [33]. In assessing the impact of BLT on sleep, both objective sleep measures, like Watch Actigraphy, and subjective sleep measures, including Sleep Disturbance Index (SDI), and Pittsburgh Sleep Quality Index (PSQI)

were employed. Agitation levels were assessed using Cohen-Mansfield Agitation Inventory (CMAI) scores, while depression was measured using Cornell Scale for Depression in Dementia (CSSD) scores, Cornell Scale for Depression in Dementia Korea Version (CSSD-K) scores, and Beck Depression Inventory. The overall mood and BPSD were evaluated through the Neuropsychiatric Inventory-Nursing Home Version (NPI-NH), the Korean Version of the Neuropsychiatric Inventory Questionnaire (KNPI-Qs), and the Interact short scale. Cognition was assessed using the Mini-Mental State Examination (MMSE). Standardised questionnaires and scales were mainly administered by nursing staff [20,31,32], except in studies where assessors were not specified [33,34] or tools like KNPI-Qs involved patient's informal caregivers [34]. Notably, in the study by Cremaşcoli, et al. [30], patients completed sleep questionnaires, specifically PSQI, under medical supervision.

Findings

Three themes were identified: 1) The impact of BLT on sleep; 2) The impact of BLT on mood and behaviour, and 3) The impact of BLT on cognition. Table 4 presents a summary of the studies of each theme.

Impact of BLT on sleep

Five studies investigated the effect of BLT on sleep outcomes such as sleep quality, rest-activity cycles, and circadian rhythm [25,30-32,34]. Objective sleep outcomes using actigraphy yielded no significant effect on sleep duration or rhythm [25,30-32,34]. However, all four studies that measured subjective sleep outcomes using proxy-rated questionnaires such as PSQI and SDI, reported significant improvements in terms of subjective sleep and total sleep time quality [30-32,34]. Variations in outcomes could be dependent on the means of assessing sleep, with significant treatment effects as measured by proxy-rated PSQI and SDI, as opposed to assessing sleep through actigraphy. Actigraphy and proxy-rated sleep measurements represented a distinct approach to measuring sleep. It is possible that actigraphy may not exhibit the same degree of sensitivity to change as proxy-rated measurements. Furthermore, low participant activity levels could

contribute to the lack of effect, as actigraphy could potentially misinterpret periods of wakefulness as sleep due to minimal movement, potentially missing subtle variations in sleep and wakefulness [32]. Nonetheless, the most used characteristics of BLT among these five studies that resulted in positive effects on sleep measures were LED lights programmed to provide varying illuminance and correlated colour temperature throughout the day, with a peak of 1000 lux, 6000 K between 10 a.m. and 3 p.m., for 24-weeks [25,32]. Overall, while BLT had a positive impact on subjective sleep outcomes, it did not show a significant effect on objective sleep measures.

Impact of BLT on mood and behaviour

Four studies investigated the impact of BLT on mood and behaviour focusing on overall BPSD and behavioural symptoms such as depression and agitation [20,30,31,34]. Firstly, Figueiro, et al. [31] reported significant improvement in agitation as measured by CMAI scores. In the four studies that measured the effect of BLT on depression, two studies reported a significant difference in CSSD scores [20,31] while the other two studies conducted by Kim, et al. [34] and Cremascoli, et al. [30] did not find any significant improvements in CSSD-K score and Beck Depression Inventory respectively. Lastly, overall BPSD was assessed using the NPI-NH and KNPI-Qs in three studies [20,30,34]. While Kolberg, et al. [20] reported significant improvements in overall BPSD, as measured by NPI-NH scores, the remaining two studies, conducted by Cremascoli, et al. [30] and Kim, et al. [34], did not report significant differences in overall BPSD. The conflicting findings may be attributed to variations in dementia severity among the participants in the four studies. The studies conducted by Figueiro, et al. [31] and Kolberg, et al. [20] focused on moderate to late-stage dementia, while the other two studies focused on early-stage dementia [30,34]. As highlighted by García-Martín, et al. [39], neuropsychiatric symptoms such as agitation and depression may be more prevalent in the later stages of dementia, which further reinforces the probability that BLT might have a more prominent impact on mood and behaviour in PwD with advanced stages of dementia. Notably, among the four studies, positive outcomes in mood and behaviour were associated with tailored BLT characteristics including LED light peaking at 1000 lux between 10 a.m. and 3 p.m. daily for 24 weeks [20], and white light 550-700 lux daily for 14-weeks [31]. Overall, BLT had a positive impact on mood and behaviour through varied outcomes. The positive effects were evident in individuals with more advanced stages of dementia but not in the early stages of dementia.

Impact of BLT on cognition

Three studies investigated the impact of BLT on cognition and consistently reported significant improvement in cognition, measured by MMSE scores [30,33,34]. Cremascoli, et al. [30] suggested that the observed improvement in cognition in PwD could have been attributed to the participants' early stage

of dementia. Yamadera, et al. [40] reported that the observed reduction in damage to the suprachiasmatic nucleus and retinogeniculate pathway during early dementia stages compared to later dementia stages may account for the cognitive improvement due to the early stage of dementia amongst the participants. However, incongruities in dementia severity, with two studies focusing on mild to moderate dementia stages [30,34] and one focusing on late-stage dementia [33], raised concerns about the generalisability and the validity of the finding, particularly in the study by Kheradmand, et al. [33]. Utilising MMSE as the primary outcome measure for cognitive assessment in the study conducted by Kheradmand, et al. [33] may potentially pose a limitation due to its sensitivity and specificity for detecting mild dementia stages, instead of advanced dementia stages [41]. Notably, the most used characteristics of BLT among these three studies involved blue-enriched light ranging from 30 to 10,000 lux for 20 to 60 minutes, daily over 2 to 4 weeks [30,34]. Overall, BLT has demonstrated positive effects on the cognitive functioning of PwD.

Discussion

Firstly, regarding the impact of BLT on sleep outcomes, studies consistently reported positive results in subjective sleep outcomes such as sleep quality and sleep disturbance, while objective sleep outcomes such as sleep-wake rhythm showed no significant positive effects across all studies [25,30-32,34]. These findings align with previous literature reporting beneficial effects on subjective sleep quality such as sleep duration as documented by improvement of PSQI scores but no difference in objective sleep measures like rest-activity rhythms [42].

In mood and behaviour assessment, particularly for overall BPSD, agitation, and depression, the results varied. Studies with advanced dementia stages showed favourable outcomes [20,31], contrasting with studies with early-stage dementia that found no significant differences [30,34]. This aligns with a previous study reporting that BLT has greater benefits in mood and behaviour for individuals in advanced stages of dementia compared to those in the early stages [43].

Cognition improved significantly in participants in all three studies that examined the impact of BLT on cognition [30,33,34]. These results were consistent with previous studies [44,45], which similarly indicated positive effects on cognitive function.

A variety of BLT characteristics was observed across all studies. However, the most used characteristic of BLT among the studies that resulted in overall positive effects on the outcome measures, particularly sleep, mood, and behaviour measures was the use of LED light for 24 weeks.

Limitations

There were several gaps found in the review of the literature. Firstly, heterogeneity among the intervention characteristics and dementia types and severity led to generally heterogeneous findings. Consequently, the overall evidence on the effectiveness of BLT for BPSD remains inconclusive. Therefore, as suggested by Nizamutdinov, et al. [46], focusing future research on moderate to late-stage dementia may yield more meaningful results, as participants with advanced stages of dementia might exhibit more noticeable improvements compared to the participants in the early stages of dementia. Additionally, employing measurement tools sensitive to changes in the later stages of dementia would enhance the accuracy of assessing BLT's impact. While MMSE is widely acknowledged as a reliable and valid measurement tool for assessing cognitive function, its greater sensitivity and specificity toward detecting mild dementia may limit its effectiveness in assessing cognitive changes in the advanced stages of dementia [47]. Future research could consider incorporating a diverse set of cognitive assessments for a more comprehensive evaluation, particularly in cases of moderate to severe dementia. Moreover, the possibility of measurement errors in light exposure duration makes it challenging to ascertain whether participants were exposed to the intended duration of light exposure. In most studies, the specific measurement of light exposure for each participant was not conducted. Instead, many of these studies relied on estimations of exposure, which were derived from standardised light meter measurements and subjective data, such as nurses' reported duration spent in the room. Therefore, to enhance the accuracy of measurement of the duration of light exposure, the development of objective light measurement tools such as wearable devices could be introduced to objectively measure the duration and amount of light to which participants are exposed.

Conclusion

This review found heterogeneity among the intervention characteristics and dementia types and severity has led to generally varied outcomes. While BLT exhibits promise as a therapeutic intervention for improving sleep, mood, behaviour, and cognition in PwD, it is strongly recommended that further research be conducted to determine the optimal implementation and efficacy of BLT across diverse populations and disease stages. By addressing these research gaps, future studies can contribute to advancing our understanding of BLT's potential as a non-pharmacological intervention for enhancing the well-being and quality of life of PwD.

References

- WHO (2022) Ageing and health. World Health Organization.
- Oh ES, Rabins PV (2019) Dementia. *Ann Intern Med* 171: ITC33-ITC48.
- WHO (2023) Dementia. World Health Organization.
- Moro E, Arabia G, Tartaglia MC, Ferretti MT (2022) Sex and gender differences in neurological disease. Academic Press is an imprint of Elsevier.
- Abraha I, Rimland JM, Trotta FM, Dell'Aquila G, Cruz-Jentoft A, et al. (2017) Systematic review of systematic reviews of non-pharmacological interventions to treat behavioural disturbances in older patients with dementia. The SENATOR- OnTop series. *BMJ Open* 7: e012759.
- Scassellati C, Ciani M, Maj C, Geroldi C, Zanetti O, et al. (2020) Behavioral and psychological symptoms of dementia (BPSD): Clinical characterization and genetic correlates in an Italian alzheimer's disease cohort. *J Pers Med* 10: 90.
- Cameron DE (1941) Studies in senile nocturnal delirium. *The Psychiatric Quarterly* 15: 47-53.
- Guu T, Aarsland D, Fyftche D (2022) Light, sleep-wake Rhythm, and behavioural and psychological symptoms of dementia in care home patients: Revisiting the Sundowning syndrome. *Int J Geriatr Psychiatry* 37.
- Canevelli M, Valletta M, Trebbastoni A, Sarli G, D'Antonio F, et al. (2016) Sundowning in dementia: Clinical relevance, pathophysiological determinants, and therapeutic approaches. *Front Med* 3: 73.
- Todd WD (2020) Potential pathways for circadian dysfunction and sundowning-related behavioral aggression in alzheimer's disease and related Dementias. *Front Neurosci* 14: 910.
- Kamoga R, Mubangizi V, Owokuhaisa J, Muwanguzi M, Natakunda S, et al. (2023) Behavioral and psychological symptoms of dementia: Prevalence, symptom severity, and caregiver distress in south-western Uganda-a quantitative cross-sectional study. *IJERPH* 20: 2336.
- Hessler JB, Schäufele M, Hendlmeier I, Junge MN, Leonhardt S, et al. (2017) Behavioural and psychological symptoms in general hospital patients with dementia, distress for nursing staff and complications in care: Results of the General Hospital Study. *Epidemiol Psychiatr Sci* 27: 278-287.
- Yang Y-y, Hsiao C-H, Chang Y-J, Ma S-C, Wang J-J (2020) Exploring dementia care competence of nurses working in Acute Care Settings. *J Clin Nurs* 31: 1972-1982.
- Yous ML, Ploeg J, Kaasalainen S, Martin LS (2019) Nurses' experiences in caring for older adults with responsive behaviors of dementia in acute care. *SAGE Open Nurs* 5: 237796081983412.
- Jennings AA, Foley T, Walsh KA, Coffey A, Browne JP, et al. (2018) General practitioners' knowledge, attitudes, and experiences of managing behavioural and psychological symptoms of dementia: A mixed-methods systematic review. *Int J Geriatr Psychiatry* 33: 1163-1176.
- Ohno Y, Kunisawa N, Shimizu S (2019) Antipsychotic treatment of behavioral and psychological symptoms of dementia (BPSD): Management of Extrapyrarnidal Side effects. *Front Pharmacol* 10: 1045.
- White N, Leurent B, Lord K, Scott S, Jones L, et al. (2017) The management of behavioural and psychological symptoms of dementia in the Acute General Medical Hospital: A longitudinal cohort study. *Int J Geriatr Psychiatry* 32: 297-305.
- Almeida OP, Ford AH (2020) Are we getting better at managing agitation in dementia? *Am J Geriatr Psychiatry* 28: 401-403.

19. Gunawardena R, Smithard DG (2019) The attitudes towards the use of restraint and restrictive intervention amongst healthcare staff on Acute Medical and frailty wards—a brief literature review. *Geriatrics* 4: 50.
20. Kolberg E, Hjetland GJ, Thun E, Pallesen S, Nordhus IH, et al. (2021) The effects of bright light treatment on affective symptoms in people with dementia: A 24-week cluster randomized controlled trial. *BMC Psychiatry* 21.
21. Jao Y-L, Wang J, Liao Y-J, Parajuli J, Berish D, et al. (2022) Effect of ambient bright light on behavioral and psychological symptoms in people with dementia: A systematic review. *Innov Aging* 6: igac018.
22. Cibeira N, Maseda A, Lorenzo-López L, Rodríguez-Villamil JL, López-López R, et al. (2020) Application of light therapy in older adults with Cognitive Impairment: A systematic review. *Geriatric Nursing* 41: 970-983.
23. Tan JSI, Cheng LJ, Chan EY, Lau Y, Lau ST (2022) Light therapy for sleep disturbances in older adults with dementia: A systematic review, meta-analysis and meta-regression. *Sleep Med* 90: 153-166.
24. Figueiro MG (2017) Light, sleep and circadian rhythms in older adults with alzheimer's disease and related dementias. *Neurodegener Dis Manag* 7: 119- 145.
25. Kolberg E, Pallesen S, Hjetland GJ, Nordhus IH, Flo-Groeneboom E (2021) The effect of bright light treatment on rest-activity rhythms in people with dementia: A 24-week cluster randomized controlled trial. *Clocks Sleep* 3: 449-464.
26. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, et al. (2021) The Prisma 2020 statement: An updated guideline for reporting systematic reviews. *BMJ* 372: n71.
27. Elsevier (2023) Mendeley reference manager. Mendeley.
28. Bromundt V, Wirz-Justice A, Boutellier M, Winter S, Haberstroh M, et al. (2019) Effects of a dawn-dusk simulation on circadian rest-activity cycles, sleep, mood and well-being in dementia patients. *Exp Gerontol* 124: 110641.
29. Cibeira N, Maseda A, Lorenzo-López L, González-Abrales I, López-López R, et al. (2021) Bright light therapy in older adults with moderate to very severe dementia: Immediate effects on behavior, mood, and physiological parameters. *Healthcare* 9: 1065.
30. Cremascoli R, Sparasci D, Giusti G, Cattaldo S, Prina E, et al. (2022) Effects of circadian phase tailored light therapy on sleep, mood, and cognition in alzheimer's disease: Preliminary findings in a pivotal study. *Front Physiol* 12: 755322.
31. Figueiro MG, Plitnick B, Roohan C, Sahin L, Kalsher M, et al. (2019) Effects of a tailored lighting intervention on sleep quality, rest-activity, mood, and behavior in older adults with alzheimer disease and related dementias: A randomized clinical trial. *J Clin Sleep Med* 15: 1757-1767.
32. Hjetland GJ, Kolberg E, Pallesen S, Thun E, Nordhus IH, et al. (2021) Ambient bright light treatment improved proxy-rated sleep but not sleep measured by actigraphy in nursing home patients with dementia: A placebo- controlled randomised trial. *BMC Geriatrics* 21: 312.
33. Kheradmand A, Donboli S, Tanjani PT, Farhadinasab A, Tabeie F, et al. (2022) Therapeutic effects of low-level laser therapy on cognitive symptoms of patients with dementia: A double-blinded randomized clinical trial. *Photobiomodul Photomed Laser Surg* 40: 632-638.
34. Kim SJ, Lee SH, Suh IB, Jang JW, Jhoo JH, et al. (2021) Positive effect of timed blue-enriched white light on sleep and cognition in patients with mild and moderate alzheimer's disease. *Sci Rep* 11: 10174.
35. Joanna Briggs Institute (2014) JBI levels of evidence.
36. Jordan Z, Munn Z, Aromataris E, Lockwood C (2015) Now that we're here, where are we? The JBI approach to evidence-based healthcare 20 years on. *Int J Evid Based Healthc* 13: 117-120.
37. Barker TH, Stone JC, Sears K, Klugar M, Tufanaru C, et al. (2023) The revised JBI critical appraisal tool for the assessment of risk of bias for randomized controlled trials. *JBI Evid Synth* 21: 494-506.
38. Hjetland GJ, Pallesen S, Thun E, Kolberg E, Nordhus IH, et al. (2020) Light interventions and sleep, Circadian, behavioral, and psychological disturbances in dementia: A systematic review of methods and outcomes. *Sleep Med Rev* 52: 101310.
39. García-Martín V, de Hoyos-Alonso MC, Ariza-Cardiel G, Delgado-Puebla R, García-Domingo P, et al. (2022) Neuropsychiatric symptoms and subsyndromes in patients with different stages of dementia in primary care follow-up (nedem project): A cross-sectional study. *BMC Geriatr* 22: 71.
40. Yamadera H, Ito T, Suzuki H, Asayama K, Ito R, et al. (2000) Effects of bright light on cognitive and sleep-wake (circadian) rhythm disturbances in alzheimer-type dementia. *Psychiatry Clin Neurosci* 54: 352-353.
41. Perneczky R, Wagenpfeil S, Komossa K, Grimmer T, Diehl J, et al. (2006) Mapping scores onto stages: Mini-mental state examination and clinical dementia rating. *Am J Geriatr Psychiatry* 14: 139-144.
42. Sloane PD, Figueiro M, Garg S, Cohen LW, Reed D, et al. (2014) Effect of home-based light treatment on persons with dementia and their caregivers. *Light Res Technol* 47: 161-176.
43. Onega LL, Pierce TW, Epperly L (2018) Bright light therapy to treat depression in individuals with mild/moderate or severe dementia. *Issues Ment Health Nurs* 39: 370-373.
44. Burns A, Allen H, Tomenson B, Duignan D, Byrne J (2009) Bright light therapy for agitation in dementia: A randomized controlled trial. *Int Psychogeriatr* 21: 711-721.
45. Riemersma-van Der Lek RF, Swaab DF, Twisk J, Hol EM, Hoogendijk WJ, et al. (2008) Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: A randomized controlled trial. *JAMA* 299: 2642-2655.
46. Nizamutdinov D, Qi X, Berman MH, Dougal G, Dayawansa S, et al. (2021) Transcranial near infrared light stimulations improve cognition in patients with dementia. *Aging Dis* 12: 954-963.
47. Baek MJ, Kim K, Park YH, Kim SY (2016) The validity and reliability of the mini- mental state examination-2 for detecting mild cognitive impairment and alzheimer's disease in a Korean population. *PLoS One* 11: e0163792.