



## ORIGINAL ARTICLE

## Impact of COVID-19 Pandemic on Elementary School Children's Sleep Hygiene

Nita Beluli Luma\*

Department of Social Work and Social Policy, Mother Theresa University, Macedonia

\*Corresponding author: Nita Beluli Luma, Department of Social Work and Social Policy, Mother Theresa University, Macedonia



### Abstract

We were challenged during the pandemic in many ways and the researchers did focus mainly on mental and physical health primarily of adults, as referred to be more at risk by the virus, yet we thought that it was essential to study how the pandemic impacted the most fragile life stages, those of the children, referring to the sleep hygiene.

Having in consideration that in our country, North Macedonia such investigations were neglected we believed that there was a need to comprehensively study the effects and outcomes of the pandemic and the safety measures on children's sleep hygiene, regarding all the consequences from the new lifestyles we got from the pandemic safety measures, such as home schooling, online learning, increased screen time, the distress etc. It is essential to study the impact of the pandemic in children referring to the sleep disturbances, having in consideration that they may trigger neurobehavioral disorders or increase the frequency of occurrence of a present, existent disease. Therefore, this study has provided us with the conclusions derived from the research we have conducted in to examine the impact of the pandemic on the probability of sleep problems in elementary school age children, highlighting the sleep patterns and specific sleep disturbances in accordance with associated factors during COVID-19 pandemic outbreak.

From the research we were able to come to conclusions that elementary school children on second grade experience higher moderate levels of these sleep disruptions: sleep anxiety; bedtime resistance; night waking and parasomnia, in comparison to 5<sup>th</sup> graders who exceed them at only one sleep subscale which is: sleep onset delay. We also came to a finding that 53% of the variance in bedtime resistance is predicted by sleep anxiety. Another conclusion brings us to an understanding that the sleep subscale of night waking can significantly predict the parasomnia sleep disorders for 45%.

This study provides us with the opportunity to conceive a general idea about children's sleep quality or sleep disruptions that will be used as guidelines on lifestyle behavior recommendations and psychotherapeutic approaches for the children.

### Keywords

Children mental health, Sleep patterns, Sleep health, Sleep disruptions, COVID-19

### Introduction

Quality and appropriate sleep are valuable for the growth and development of children. Referring to the latest findings and studies which have been conducted to research about the impact of current pandemic on children's sleep hygiene and their overall wellbeing have found frequent alterations in sleep habits; increased total sleep duration and negative impact on sleep quality.

Sleep disorders bring many problems along with them. Therefore, many correlated risk factors of distress have been investigated as independent variables, where many authors have come to an understanding that there is an increase in sleep disorders that significantly impact the quality of children's life and mental wellbeing [1].

The psychological distresses due to the pandemic outbreak have become part of all major stages of human life circle, without excluding the children. Especially during the lockdown, as a safety measure, which confined the children to their homes for an extended period of time, with schools remaining closed and students only allowed to follow online lessons, in order

to reduce contagion and pressure on the healthcare system, there have always been concerns that because of the prolonged home confinement during a disease outbreak may affect physical and mental health of children [2], and not only them.

For the sake of children's wellbeing and in order to maintain their potential at doing their assessments, it is crucial for them that they have a good quality of sleep at night. During the home confinement the sleep issues emerge, which only triggers an increase in stress and anxiety levels and which may possibly bring a decrease in the overall quality of their life.

The loss of social contacts; the reduction of physical activity; the need to play with their school-mates in order to achieve their social development skills; the lack of sunlight exposure which may as well intrude with the flexibility in wake/sleep time, socio-emotional struggles due to changes in their family dynamic and overall mood among the family members due to various factors, the increase of screen-time and the changes in their eating habits, and many more to count, are enough reasons to doubt that the children in our place are facing with not so common sleeping problems. These changes can not only impact daily activities as well as the sleep/wake pattern and circadian rhythmicity [3].

Having in mind that sleep disorders may trigger many other psychological issues along with them, and that the quality of sleep is very important for children's growth and development, we find it very crucial to study the prevalence of sleep disruptions during the pandemic COVID-19 among children.

## Methodology

### Purpose

The purpose of the present study is to provide detailed data of the impact of COVID-19 pandemic outbreak on children sleep patterns and sleep disturbances, as well as highlighting the importance of the link between sleep health and family related factors [4,5].

With this survey, we aimed to evaluate the effects of COVID-19 pandemic on sleep quality as an indicator of psychological well-being among children living in North Macedonia. Secondly, we aimed to identify potential familial, socioeconomic, and personal risk factors for their occurrence [6,7].

### Research methods

We have selected the participants of this study by the so called cluster sampling" in the city of Struga, in a public elementary school, North Macedonia. The survey included 85 respondents divided by their school year (2<sup>nd</sup> and 5<sup>th</sup> grade) [8-10].

For the assessment of the sleep patterns and disturbances among children we have used the Children Sleep Habits Questionnaire which is a parent

- rated questionnaire that evaluates common pediatric sleep difficulties. For data collection we have used the 'snowball' sampling technique, which is an online software platform, the one we used is called Survey Planet. The data were analyzed with the SPSS statistical software version 22 [11-13].

## Hypothesis

*H1. 2<sup>nd</sup> grade children experience higher levels of sleep disruptions than 5<sup>th</sup> graders in all sleep subscales.*

*H2. Younger school age children experience more sleep disturbances and anxiety than older one's.*

*H3. Sleep anxiety and bedtime resistance positively correlate with one another.*

*H4. Sleep anxiety impacts the onset of bedtime resistance behaviors in elementary school children.*

*H5. Children who experience night waking are more predisposed to experience parasomnia issues.*

## Findings and results

Regarding the data that we have analyzed, we have been able to come to these findings which indicate that 2<sup>nd</sup> graders experience higher moderate levels of sleep disruptions, while high levels of these disruptions weren't met in representative percentage to be mentioned [14,15].

We have come to conclusion that children at age 7-8 that experience moderate levels of sleep anxiety are (% = 48.8) of our population, while only (% = 27.3) from 5<sup>th</sup> graders are experiencing moderate levels of sleep anxiety. Bedtime resistance is also more present at 2<sup>nd</sup> graders (% = 43.90) whose percentage of those who exhibit these issues in moderate levels exceed those of 5<sup>th</sup> graders (% = 27.27) significantly. Night waking issues are expressed in a moderate level at younger pupils in (%29.27), while 10-11 years-old pupils (%18.18) exhibit such sleep issues. For parasomnia disorders we have found out that such symptoms are detected at (% = 29.27) of 2<sup>nd</sup> graders and in (% = 13.64) of 5<sup>th</sup> graders. Daytime sleepiness disorder hasn't been shown to be a concern in our population, since more than 92% of the respondents haven't met it's symptoms in their everyday life. The only sleep disruption in which 5<sup>th</sup> graders exceed with percentage of the sleep issues presence in moderate level, the 2<sup>nd</sup> graders, is sleep onset delay, on which 5<sup>th</sup> graders achieve (% = 38.64) and 2<sup>nd</sup> graders have (% = 26.83) [15,16].

Through the implementation of this study we have been able to come to a finding that sleep anxiety and bedtime resistance are positively correlated with one another. The correlation is ( $r = 0.731$ ) which indicates strong correlation between these two variables, meaning that with an increase in sleep anxiety the children's probability of bedtime resistance will also grow, and vice versa (Table 1) [17,18].

**Table 1:** Pearson correlation of sleep anxiety and bedtime resistance.

Correlations			
		Sleep anxiety	Bedtime resistance
Sleep anxiety	Pearson Correlation	1	0.731**
	Sig. (2-tailed)		0
	N	85	85
Bedtime resistance	Pearson Correlation	0.731**	1
	Sig. (2-tailed)	0	
	N	85	85

\*\* : Correlation is significant at the 0.01 level (2-tailed).

**Table 2:** Regression test.

Model summary <sup>b</sup>				
Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	0.731 <sup>a</sup>	0.534	0.528	2.247

<sup>a</sup>: Predictors: (Constant), Sleep Anxiety; <sup>b</sup>: Dependent Variable: Bedtime resistance

**Table 3:** Anova test.

ANOVA <sup>a</sup>						
Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	480.192	1	480.192	95.109	0.000 <sup>b</sup>
	Residual	419.055	83	5.049		
	Total	899.247	84			

<sup>a</sup>: Dependent variable: Bedtime resistance; <sup>b</sup>: Predictors: (Constant), Sleep anxiety

**Table 4:** Test of equality of covariance matrices.

Box's Test of Equality of Covariance Matrices <sup>a</sup>	
Box's M	1.665
F	0.54
df1	3
df2	1483448.696
Sig.	0.655

Tests the null hypothesis that the observed covariance matrices of the dependent variables are equal across groups.

<sup>a</sup>: Design: Intercept + Q1

Regression test has confirmed our hypothesis that sleep anxiety does influence the onset of bedtime resistance issues. Through our findings we were able to come to the following findings:

The R value represents the simple correlation and is  $R = 0.73$  which indicates a high degree of correlation, while looking at the adjusted R square data, we come to an understanding that  $R^2 (0.53 \times 100) = 53\%$  of the variance in bedtime resistance is predicted by sleep anxiety (Table 2).

This table (Table 3) indicates that the regression model predicts the dependent variable significantly well. The model is significant  $F (1.83) = 96.11$ , ( $p = 0.00 < 0.01$ ), which indicates that, overall, the regression model statistically significantly predicts the outcome variable. Sleep anxiety accounted for 53%

of the explained variability in bedtime resistance. The regression equation was: predicted bedtime resistance =  $2.43 + 9.21 \times (\text{sleep anxiety})$  [19,20].

The table of the test of equality if covariance matrices (Table 4), provides evidence that equal variance assumption is satisfied since ( $p = 0.655 > 0.01$ ).

There was a significant difference between younger (7-8 y.o) and older pupils (10-11 y.o) when considered jointly on the variables of sleep issues in general and sleep anxiety, Wilks  $\Lambda = 0.85$ ,  $F = 7.36$ ,  $p = 0.01$ , partial  $\eta^2 = 0.15$ . A separate ANOVA was conducted on each dependent variable, with each ANOVA evaluated at an alpha level of 0.025. There was a significant difference between 2<sup>nd</sup> and 5<sup>th</sup> graders on general sleep disruptions  $F (1.83) = 10.98$ , partial  $\eta^2 = 0.12$ , with 2<sup>nd</sup> graders ( $M = 45.54$ ) scoring higher than 5<sup>th</sup> graders ( $M = 39.98$ ). Also there was a significant difference between 2<sup>nd</sup> and 5<sup>th</sup> graders on general on sleep anxiety  $F (1.83) = 14.78$ , partial  $\eta^2 = .0.15$  with means with 2<sup>nd</sup> graders ( $M = 24.17$ ) scoring higher than 5<sup>th</sup> graders ( $M = 19.52$ ) (Table 5 and Table 6).

A linear regression established that night waking could significantly predict parasomnia issues  $F (1.83) = 70.57$ ,  $p = 0.001 < 0.01$  and night waking accounted for 45% of the explained variability in parasomnia sleep disorders (Table 7).

These findings confirm our hypothesis that "children

**Table 5:** MANOVA Test.

Multivariate tests <sup>a</sup>							
Effect		Value	F	Hypothesis df	Error df	Sig.	Partial Eta Squared
Intercept	Pillai's Trace	0.974	1540.112 <sup>b</sup>	2	82	0	0.974
	Wilks' Lambda	0.026	1540.112 <sup>b</sup>	2	82	0	0.974
	Hotelling's Trace	37.564	1540.112 <sup>b</sup>	2	82	0	0.974
	Roy's Largest Root	37.564	1540.112 <sup>b</sup>	2	82	0	0.974
School grade	Pillai's Trace	0.152	7.365 <sup>b</sup>	2	82	0.001	0.152
	<b>Wilks' Lambda</b>	<b>0.848</b>	<b>7.365<sup>b</sup></b>	<b>2</b>	<b>82</b>	<b>0.001</b>	<b>0.152</b>
	Hotelling's Trace	0.18	7.365 <sup>b</sup>	2	82	0.001	0.152
	Roy's Largest Root	0.18	7.365 <sup>b</sup>	2	82	0.001	0.152

<sup>a</sup>: Design: Intercept + Q1; <sup>b</sup>: Exact statistic

**Table 6:** Test of between subjects when considered jointly.

Tests of Between-Subjects Effects							
Source	Dependent variable	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Sq
Corrected Model	Total sleep	655.933 <sup>a</sup>	1	655.933	10.978	0.001	0.117
	Sleep anxiety	458.512 <sup>b</sup>	1	458.512	14.78	0	0.151
Intercept	Total sleep	155199.6	1	155199.6	2597.524	0	0.969
	Sleep anxiety	40518.23	1	40518.23	1306.135	0	0.94
<b>School year</b>	<b>Total sleep</b>	<b>655.933</b>	<b>1</b>	<b>655.933</b>	<b>10.978</b>	<b>0.001</b>	<b>0.117</b>
	<b>Sleep anxiety</b>	<b>458.512</b>	<b>1</b>	<b>458.512</b>	<b>14.78</b>	<b>0</b>	<b>0.151</b>
Error	Total sleep	4959.172	83	59.749			
	Sleep anxiety	2574.782	83	31.021			
Total	Total sleep	160296	85				
	Sleep anxiety	43298	85				
Corrected Total	Total sleep	5615.106	84				
	Sleep anxiety	3033.294	84				

<sup>a</sup>: R Squared = 0.117 (Adjusted R Squared = 0.106); <sup>b</sup>: R Squared = 0.151 (Adjusted R Squared = 0.141)

**Table 7:** Linear regression.

Model Summary <sup>b</sup>				
Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	0.678a	0.46	0.453	2.12

<sup>a</sup>: Predictors: (Constant), Night awakenings; <sup>b</sup>: Dependent variable: Parasomnia total

Coefficients <sup>a</sup>								
		Unstandardized Coefficients		Standardized Coeffi	95.0% Confidence Interval for B			
Model		B	Std. Error	Beta	T	Sig.	Lower Bound	Upper Bound
1	(Constant)	3.492	0.707		4.942	0	2.087	4.897
	Night awakenings	0.912	0.109	0.678	8.402	0	0.696	1.128

<sup>a</sup>: Dependent variable: Parasomnia total

who experience night waking are more predisposed to experience parasomnia sleep issues”.

## Conclusions

Regarding our findings we have ended up with the

following conclusions:

Second graders experience higher moderate levels of these sleep disruptions: Sleep anxiety; bedtime resistance; night waking and parasomnia, in comparison to 5<sup>th</sup> graders who exceed them at only

one sleep subscale which is: sleep onset delay. On the subscale of daytime sleepiness disorder, weren't found any significant differences between the respondents because the majority of each group of respondents didn't manifest any symptoms correlated to this sleep disorder.

We came to a conclusion that sleep anxiety and bedtime resistance are positively correlated with each other, meaning that the pupils that experience higher levels of anxiety are those who are more highly predisposed to exhibit bedtime resistance as well.

Throughout the implementation of the linear regression test we came to a finding that 53% of the variance in bedtime resistance is predicted by sleep anxiety.

We also came to an interesting finding through the MANOVA tests, that there was a significant difference between 2<sup>nd</sup> graders (7-8 y.o) and pupils on 5<sup>th</sup> grade (10-11 y.o) when considered jointly on the variables of sleep issues in general and sleep anxiety, having the 2<sup>nd</sup> graders in advantage, scoring higher in both mentioned variables.

And the last but not the least conclusion brings us to an understanding that the sleep subscale of night waking can significantly predict the parasomnia sleep disorders for 45%.

In summary, our research study findings have highlighted the importance of children sleep health during the pandemic outbreak. I think that future studies should include assessment of other factors that could be related to sleep disruptions such as health concerns, academic performance during the period through e-learning, the family financial stability, social anxiety due to social isolation etc. in order to further explore and contribute in children's mental health.

## References

- Dondi A, Fetta A, Lenzi J, Morigi F, Candela E, et al. (2021) Sleep disorders reveal distress among children and adolescents during the Covid-19 first wave: Results of a large web-based Italian survey. *Ital J Pediatr* 47: 130.
- Wang C, Pan R, Wan X, Tan Y, Xu L, et al. (2020) Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health* 17: 1729.
- Altena E, Baglioni C, Espie CA, Ellis J, Gavriloff D, et al. (2020) Dealing with sleep problems during home confinement due to the COVID-19 outbreak: Practical recommendations from a task force of the European CBT-I Academy. *J Sleep Res* 29: e13052.
- Becker SP, Gregory AM (2020) Editorial Perspective: Perils and promise for child and adolescent sleep and associated psychopathology during the COVID-19 pandemic. *J Child Psychol Psychiatry* 61: 757-759.
- Bhargava S (2011) Diagnosis and management of common sleep problems in children. *Pediatr Rev* 32: 91-98.
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, et al. (2020) The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *Lancet* 395: 912-920.
- Carter KA, Hathaway NE, Lettieri CF (2014) Common sleep disorders in children. *Am Fam Physician* 89: 368-377.
- Dellagiulia A, Lionetti F, Fasolo M, Verderame C, Sperati A, et al. (2020) Early impact of COVID-19 lockdown on children's sleep: A 4-week longitudinal study. *J Clin Sleep Med* 16: 1639-1640.
- Golberstein E, Wen H, Miller BF (2021) Coronavirus disease 2019 and effects of school closure for children and their families-reply. *JAMA Pediatr* 175: 211-212.
- Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R (2020) Effects of Covid-19 lockdown on mental health and sleep disturbances in Italy. *Int J Environ Res Public Health* 17: 4779.
- Kieckhefer GM, Ward TM, Tsai S-Y, Lentz MJ (2008) Nighttime sleep and daytime nap patterns in school age children with and without asthma. *J Dev Behav Pediatr* 29: 338-344.
- Liu Z, Tang H, Jin Q, Wang G, Yang Z, et al. (2021) Sleep of preschoolers during the coronavirus disease 2019 (COVID-19) outbreak. *J Sleep Res* 30: e13142.
- Luma NB (2022) Sleep patterns and sleep disturbances in school-age children amid covid-19 pandemic outbreak. *European Journal of Psychological Research* 9: 1-10.
- Messner AH, Pelayo R (2000) Pediatric sleep-related breathing disorders. *Am J Otolaryngol* 21: 98-107.
- Moore SA, Faulkner G, Rhodes RE, Brussoni M, Chulak-Bozzer T, et al. (2020) Impact of the COVID-19 virus outbreak on movement and play behaviours of Canadian children and youth: A national survey. *International Journal of Behavioral Nutrition and Physical Activity* 17: 85.
- Owens J (2007) Classification and epidemiology of childhood sleep disorders. *Sleep Medicine Clinics* 2: 353-361.
- Pianosi P (1999) Sleep disorders in children and adolescents. *Advances in Psychiatric Treatment*.
- Sadeh A, Gruber R, Raviv A (2000) Sleep patterns and sleep disruptions in school-age children. *Dev Psychol* 36: 291-301.
- Sprang G, Silman M (2013) Posttraumatic stress disorder in parents and youth after health-related disasters. *Disaster Med Public Health Prep* 7: 105-110.
- Uema SFH, Vidal MVR, Fujita R, Moreira G, Pignatari SSN (2006) Behavioral evaluation in children with obstructive sleep disorders. *Braz J Otorhinolaryngol* 72: 120-122.