



## RESEARCH ARTICLE

## Incidence of Mouth Cancer in Iraq

Hala Nadhim Kadhim, BSc, MSc<sup>1</sup> and Nadham Kadham Mahdi, MSc, PhD<sup>2\*</sup> 

<sup>1</sup>University of Al-Watania, Thi-Qar, Iraq

<sup>2</sup>Department of Microbiology, College of Medicine, University of Basrah, Basrah, Iraq

\*Corresponding author: Dr. Nadham K. Mahdi, MSc, PhD, Department of Microbiology, College of Medicine, University of Basrah, Basrah, Central Post Office-42001, P. O. Box 1565, Basrah, Iraq



### Abstract

The incidence of mouth cancers among Iraqi people is reviewed from 1973-2021 in different Iraqi provinces. The incidence rates are also illustrated in relation to age, sex, site of cancer and year of registration. Incidence rate differs widely in relation to locality, age, sex and race. The risk factors for the incidence can be attributed to tobacco and/or alcohol consumption. The relationship between diet and nutrition to the risk of cancer development has been established by several epidemiological and laboratory studies. They indicated that low intake of fruits and vegetables predisposes to increased risk of cancer development. Other risk factors include genetic, sun exposure, mate drinking (tea-like beverage), viral infection, fungal infection and chronic trauma.

Adults and children of both sexes are infected. All authors in Iraq indicated that tongue is the most common site for mouth cancer among Iraqi people. Implementing a national control program should include a primary health care, health education, well-balanced diet, environmental sanitation and health education to stress the important of the hazard of tobacco and alcohol. The knowledge about mouth cancers considerably increased when the subjects received information from their dentists.

### Keywords

Epidemiology, Incidence, Mouth cancer

### Introduction

Mouth cancers are distributed worldwide. They are responsible for millions of morbidity and mortality. Thus, it is a public health problem in many parts of the world. In addition, cancer might be undetectable and unrecorded in many countries. Oral squamous cell carcinoma is the commonest oral malignancy. It is incident in middle-aged and elderly, although it has been

recorded in young adults [1]. An estimated 26, 3000 new mouth cancer cases were reported annually over the world which accounts 2.1% of all new cases were recorded [2]. Socio-cultural behaviour for population has an important role in the geographical distribution of the disease [3]. Incidence rate differs widely in relation to locality, age, sex and race. Few studies were carried out in Arab world in relation to the incidence of mouth cancer including Kuwait [4], United Arab Emirates [5], Sudan [6], Saudi Arabia [7] and Jordan [8] as well as in Iraq (The present review).

An increase in incidence was noticed in United Kingdom [9], Netherland and Denmark [10], India, Pakistan and Bangladesh [11]. In contrast, a decrease in the incidence have reported in USA, Italy, Hong Kong, France, Germany and Australia [12].

In Iraq, mouth cancer represents 4.5% of all malignant cancers as recorded by Iraq cancer registry [13], the present review aimed to determine the analysis of Iraqi mouth cancers data according to age, sex, year of registration and cancer site over time.

### Results and Discussion

The incidence rates of mouth cancer are reviewed according to types of patients including age and sex in different Iraqi provinces (Table 1). The incidence is varying from one region to another.

Fuoad, et al., [14] result has indicated a slight increase in the rate of mouth cancer over time (14.5%) compared to a five-year (2004-2009) retrospective study (12.3%) in Sulaimania [15] and another previous study in Baghdad during 1991-2000 (9.97%) [16] as well as in

**Table 1:** Distribution of mouth cancer in relation to province, years of registration, age and gender.

Ref. No.	Province	Years of incidence	Age (Years)	Sex	
				Male No. (%)	Female No. (%)
Perriman [18]	Baghdad	1967-1972	11-80	13 (56.5%)	10 (43.5%)
Al-Niimi [19]	Mosul	1995-1999	2.5-90	64 (53.8%)	55 (46.2%)
Talabani, et al. [34]	Sulaimania	1995-2004	51-60	44 (60.3%)	29 (39.7%)
Museedi & Younis [22]	Baghdad	2000-2008	-	1035 (57.9%)	752 (42.1%)
Taha & Younis [20]	Baghdad	2001-2013	< 20-70	976 (58.7%)	688 (41.3%)
Al-Kawaz [35]	Baghdad	2003-2006	51-60	80 (62%)	49 (38%)
Khudier [15]	Sulaimania	2004-2008	50-70	M:F - 1.2:1	
Fuoad, et al. [14]	Sulaimani	2008-2019	1-90	180 (53.4%)	157 (46.6%)
Aljazaeri, et al. [21]	Basrah	2012-2017	1-85	283 (50.54%)	277 (49.46%)
Al-Mahfoud, et al. [23]	Basrah	2015-2016	11-80	31 (79.5%)	8 (25.5%)
Alshami, et al. [36]	Baghdad	2019	18-80	128 (40.3%)	190 (59.7%)

**Table 2:** Incidence of mouth cancers and their types in the Iraqi provinces.

Ref. No.	No. Examined	No. (%) incidence of oral cancer
Perriman [18]	202	Palat-19 (11.3%), tongue-38 (22.5%), lips-50 (29.5%), maxilla-21 (12.3%), floor of the mouth-26 (15.4%), buccal commissure-47 (26.6%), retromolar regions-14 (8.2%).
Al-Niimi [19]	119	Palat-14 (11.7%), tongue-28 (23.5%), lips-49 (41.1%), floor of the mouth-9 (7.5%), buccal mucosa -8 (6.7).
Talabani, et al. [34]	73	Tongue-16 (21.9%), unspecified-57 (78.1%).
Museedi & Younis [22]	1787	Tongue (0.47%), lips (0.3%), salivary gland (0.195).
Taha & Younis [20]	1664	Lip-382 (23%), tongue -711 (42.75%), Gingiva-83 (5.09%), Floor of the mouth-98 (5.9%), palate 72-(4.3%), unspecified- 318 (19.1%).
Al-Kawaz [35]	129	Tongue-72 (55.8%), floor of the mouth-5 (3.9%), unspecified-52 (40.3%).
Khudier [15]	82	Lower lip (39%), tongue (20%), salivary gland (19.6%).
Fuoad, et al. [14]	302	Palat-33 (10.9%), tongue- 50 (16.5%), lip-39 (12.9%), oropharynx-62 (20.5%), maxilla-45 (14.9%), jaw-38 (12.6%), salivary gland-35 (11.7%).
Aljazaeri, et al. [21]	560	Tongue-101 (18%), buccal mucosa-89 (15.9%), salivary gland- 34 (6.1%), gingival-88 (15.7%).
Al-Mahfoud, et al. [23]	21	Palat-3 (14.3%), tongue-5 (23.8%), cheek-3 (14.28%), lips-2 (9.52%), oropharynx-3 (14.3%), post nasal space-1 (4.76%), salivary gland-1 (4.76%), jaw-1 (4.76%).
Alshami, et al. [36]	318	Unspecified

the present review (Table 1 and Table 2). This can be associated to pollution, recurrent explosions and wars in Iraq. Furthermore, it might be related to the practices of tobacco smoking and drinking alcohol in both men and women which increased the risk of lip cancer worldwide [17], including Iraq [15,18-20]. All authors in Iraq indicated that tongue is the most common site for mouth cancer among Iraqi people (Table 2) which is in agreement with studies in Europe and USA [11]. Buccal mucosa is the most common site in Mosul [19], Basrah [21] as well as among Asian populations due to betel quid/tobacco chewing habits [11]. Cheek was the dominant site in women, while lip, face and mouth floor were observed among men. Lip, tongue, gum, palate found among males than females [18-20,22,23].

Mouth cancer affected males more than females

(Table 1). This is in agreement with most similar studies in countries around the world where the ratio is 1.5:1 [11]. The incidence can be explained by the contributions of smoking, alcohol or hormonal factors as well as exposure to radiation due to men's job which lead to incidence of lip cancer [17]. More than 80% of mouth cancers can be attributed to tobacco and/or alcohol consumption [24]. The relationship between diet and nutrition to the risk of cancer development has been established by several epidemiological studies [25]. They indicated that low intake of fruits and vegetables predisposes to increased risk of cancer development. Certain food as processed meats, cakes, desserts, butter, eggs, soups, red meat, salted meat, cheese, pasta or rice and corn bread [26]. More frequent consumption of fruit and vegetables, particularly of carrots, fresh tomatoes and

green peppers were associated with reduced risk of oral and pharyngeal cancer [27] as well as other food as fish, vegetable oil, bread, cereals, protein, fat, fresh meat, chicken, liver, shrimp and fiber [28]. Other risk factors include genetic [29], sun exposure [30], mate drinking (tea-like beverage) [27], viral infection [31], fungal infection [32] and chronic trauma [33].

Age-group 61-70 years was highly affected as stated by the Iraqi Cancer Registry [22], in Sulaimani [15] and in Basrah [21] from 1). Age ranged 41-60 years showed tongue cancer more other types [22].

## Conclusion

Mouth cancer is incident among all types of communities and population samples from both urban and rural regions of Iraq. Adults and children of both sexes are infected. It is an important disease and remain challenge to the dentist and since the early diagnosis of oral cancer is vital in the treatment and prognosis of mouth cancer. Dentist should continue to be encouraged to perform mouth cancer examination for all patients.

Therefore, due to lack of effective chemotherapy or vaccine against cancer, an urgent and efficient preventive and control measures is essential. Implementing a national control program should include a primary health care, health education, well-balanced diet, environmental sanitation and health education to stress the important of the hazard of tobacco and alcohol. Well trained health workers chosen from the same community are valuable in the diagnosis and treatment especially in rural areas and far villages in the country. The knowledge about mouth cancers considerably increased when the subjects received information from their dentists.

## Committee Approval

Ethics committee approval was not requested for this study.

## Conflict of Interest

None declared.

## Funding

Self-funding.

## Authorship Contributions

Authors were involved in collecting data, analysis and righting up.

## References

- Alvarez-Amézaga J, Barbier-Herrero L, Pijoan-Zubizarreta JI, Martín-Rodríguez JC, Romo-Simón L, et al. (2007) Diagnostic efficacy of sentinel node biopsy in oral squamous cell carcinoma. Cohort study and meta-analysis. *Med Oral Patol Oral Cir Bucal* 12: 235-243.
- Ferlay J, Shin HR, Bray F, Forman D, Mathers C, et al. (2010) GLOBOCAN 2008, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 10. Lyon, France: International Agency for Research on Cancer.
- Enwonwu CO, Meeks VI (1995) Bionutrition and oral cancer in humans. *Crit Rev Oral Biol Med* 6: 5-17.
- Joseph BK, Ali MA, Dashti H, Sundaram DB (2019) Analysis of oral and maxillofacial pathology lesions over an 18-year period diagnosed at Kuwait University. *J Invest Clin Dent* 10: e12432.
- Anis R, Gaballah K (2013) Oral cancer in the UAE: A multicenter, retrospective study. *Libyan J Med* 8: 21782.
- Osman T, Satti A, Bøe O, Yang Y, Ibrahim S, et al. (2010) Pattern of malignant tumors registered at a referral oral and maxillofacial hospital in Sudan during 2006 and 2007. *J Cancer Res Ther* 6: 473-477.
- AlHindi NA, Sindi AM, Binmadi NO, Elias WY (2019) Retrospective study of oral and maxillofacial pathology lesions diagnosed at the Faculty of Dentistry, King Abdulaziz University. *Clin Cosmet Investig Dent* 1: 45-52.
- Telfah H, Hammouri E (2014) Malignant tumors of the oral cavity among Jordanians. *JRMS* 21: 6-13.
- Conway DI, Stockton DL, Warnakulasuriya KA, Ogden G, Macpherson LM (2006) Incidence of oral and oropharyngeal cancer in United Kingdom (1990-1999) - recent trends and regional variation. *Oral Oncol* 42: 586-592.
- Visser BJO, Leemans CR (2009) Oral and oropharyngeal cancer in The Braakhuis, Netherlands between 1989 and 2006 Increasing incidence but not in young adults. *Oral Oncol* 45: 85-89.
- Warnakulasuriya S (2009) Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol* 45: 309-316.
- Yako-Suketomo H, Matsuda T (2010) Comparison of time Trends in lip, oral cavity and pharynx cancer mortality (1990-2006) between countries based on the WHO mortality database. *Japan J Clin Oncol* 40: 1118-1119.
- TalabaniNGh, Al-Rawi NH (2002) Squamous cell carcinoma of oral cavity: A clinico-pathological analysis of 1425 cases from Iraq. *Al-Rafidan Dent J* 1: 1-6.
- Fuoad SAA, Mohammad DN, Hamied MA, Garib BT (2021) Oro-facial malignancy in north of Iraq: a retrospective study of biopsied cases. *BMC Oral Health* 21: 1-18.
- Khudier H (2012) Malignant oral lesions in Sulaimania governorate (5 years): Retrospective study. *Diyala J Med* 2: 13-20.
- Al-Reyahi AB (2004) Retrospective analysis of malignant oral lesions for 1534 patients in Iraq during the period (1991-2000). M.Sc. Thesis, College of Dentistry, University of Baghdad.
- Sugerman PB (2000) Prevention of oral cancer current medical literature. *J Ear Nose Throat* 1: 57-61.
- Perriman A (1973) Oral cancer in Iraq: Analysis of 202 cases. *British Journal of Oral Surgery* 11: 146-151.
- Al-Niaini AI (2006) Oral malignant lesions in a sample of patients in the north of Iraq (Retrospective study). *Al-Rafidain Dent J* 6: 176-180.
- Taha I, Younis W (2015) Clinicopathological analysis of oral squamous cell carcinoma in Iraq during period (2001-2013). *J Baghdad Coll Dentistry* 27: 58-65.
- Aljazeerai SAW, Al Qudsi GH, Jaber HK, Al Elwi WM, Haddad SS, et al. (2020) Biopsy records to the oral lesions in Basrah between 2012-2017. *Journal Oral Medicine Oral Surg Oral Pathol Oral Radiol* 6: 74-80.
- Museed OS, Younis WH (2014) Oral cancer trends in Iraq from 2000-2008. *The Saudi Journal for Dental Research* 5: 41-47.

23. Al-mahfoud MM, AlSaimary IE, Al-shawi A (2017) Occurrence of oral and oropharyngeal squamous cell carcinoma among patients in Basrah city. *Iraqi J Biotechnol* 16: 255-261.
24. Radoi L, Paget-Bailly S, Cyr D, Papadopoulos A, Guida F (2013) Tobacco Smoking, Alcohol Drinking and Risk of Oral Cavity Cancer by Subsite: Results of a French Population-Based Case-Control Study, the ICARE Study. *Eur J Cancer Prev* 22: 268-276.
25. De Stefani E, Ronco A, Mendilaharsu M, Deneo-Pellegrini H (2000) Diet and risk of cancer of the upper aerodigestivetract-II. *Nutrients. Oral Oncol* 35: 22-26.
26. Bernzweig E, Payne JB, Reinhardt RA, Dyer JK, Patil KD (2000) Nicotine and smokeless tobacco effects on gingival and peripheral blood mononuclear cells. *J Clin Periodontal* 25: 246-252.
27. Dikshit RP, Kanhere S (2000) Tobacco habits and risk of lung, oropharyngeal and oral cavity cancer: A population-based case-control study in Bhopal, India. *Int J Epidemiol* 29: 609-614.
28. Jeng JH, Chang MC, Hahn LJ (2001) Role of areca nut in betel quid-associated chemical carcinogenesis: Current awareness and future perspectives. *Oral Oncol* 37: 477-492.
29. Lu D, Yu X, Du Y (2016) Meta-Analyses of the effect of cytochrome P450 2E1 gene polymorphism on the risk of head and neck cancer. *Molecular Biol Rep* 38: 2409-2416.
30. Johnson NW, Amarasinghe HK (2016) Epidemiology and aetiology of head and neck cancers. Bernier J (ed). *Head and Neck Cancer: Multimodality Management*. Springer 1: 1-58.
31. James PD, Ellis IO (2000) Malignant epithelial tumors associated with autoimmune sialadenitis. *J Clin Pathol* 39: 497-502.
32. Zain RB, Fukano F, Razak IA, Shanmuhasuntharam P, Ikeda N, et al. (2000) Risk factors for oral leukoplakia: A comparison of two ethnic groups in Malaysia. *Oral Oncol India: Macmillan* 37: 207-210.
33. Piemonte ED, Lazos JP, Brunotto M (2010) Relationship between chronic trauma of the oral mucosa, oral potentially malignant disorders and oral cancer. *J Oral Pathol Med* 39: 513-157.
34. Talabani NG, Ahmed KM, Faraj FH (2010) Oral Cancer in Sulaimani: A Clinicopathological Study. (JZS) *J Zankoy Sulaimania* 13 Part A: 1-3.
35. Al-Kawaz AB (2010) Oral squamous cell carcinoma in Iraq: clinical analysis. *Mustansiria Dental J* 7: 100-105.
36. Alshami ML, Abdulbaqi HR, Abdulkareem AA (2019) Awareness and knowledge of oral cancer in the city of Baghdad, Iraq: A questionnaire-Based survey. *J Stoma* 72: 263-268.