



RESEARCH ARTICLE

Impact of a Novel Training Approach on Body Composition in the Elderly

Matthew C Scott¹, Jason D. Allen^{2,3}, Neil M. Johannsen^{1,4}, Daniel P. Credeur⁵, Conrad P. Earnest⁶, Timothy S. Church⁴, Eric Ravussin⁴, William E. Kraus³ and Michael A. Welsch^{7*}

¹School of Kinesiology, Louisiana State University, USA

²Department of Kinesiology, University of Virginia, USA

³Duke Molecular Physiology Institute, Duke University School of Medicine, USA

⁴Pennington Biomedical Research Center, USA

⁵School of Kinesiology and Nutrition, University of Southern Mississippi, USA

⁶Department of Health and Kinesiology, Texas A&M University, USA

⁷School of Population Health, University of Mississippi Medical Center, USA

*Corresponding author: Michael Andrew Welsch, Ph.D., FACSM, Professor of Population Health Science, The John D. Bower School of Population Health, University of Mississippi Medical Center, 2500 N. State St. Jackson, MS 39216, USA, Tel: (601)-815-9553



Abstract

Background/Purpose: Deterioration of body composition with advancing age is related to functional decline. This study examined the effects of a novel progressive exercise training program on body composition in individuals > 70 y, at risk of losing functional independence.

Methods: Fifty participants (33 females, 75 ± 4.5 y) performed 45-60 min sessions, 3 times per week, for a total of 12 weeks. Training consisted of 2 distinct phases. For Phase I (First 4 weeks) participants were randomized to aerobic training consisting of walking or cycling at 50% of heart rate reserve, or PRIME (Peripheral Remodeling through Intermittent Muscular Exercise) designed to stimulate the major muscle groups (8 targeted exercises) at 40%-50% of maximal voluntary capacity, for a duration of 3-6 min, or 45 to 90 contractions. Phase I was followed by 8 weeks of combined aerobic and resistance training using established exercise guidelines (phase II). Body composition (Fat Free Mass, Fat Mass, body fat %, Skeletal Muscle Index, bone mineral content and density) were measured using Dual X-ray Absorptiometry, before (V1), and after Phase I (V2) and II (V3).

Results: Compared to V1, V3 values for Body Fat% (p = 0.005) decreased, and Fat Free Mass (p = 0.03) and Skeletal Muscle Index (p = 0.05) increased, in both groups. Changes

in Bone Mineral Content and Bone Mineral Density were significantly greater for PRIME compared to the aerobically trained group (p < 0.05).

Conclusion: Both groups improved body fat percentage, increased SMI, and BMC and BMD, in specific sites measured. Uniquely, participants whom initiated training with PRIME experienced greater improvements in BMC and BMD. These data support a paradigm shift in the order of exercise training, for older adults at risk of losing independence, with initial emphasis on a moderate to high intensity localized exercise stimulus, rather than moderate to high intensity systemic exercise.

Keywords

Exercise, Skeletal muscle index, Bone mineral density, Aging

Introduction

Aging is associated with significant changes in body composition. Evidence from the Baltimore Longitudinal Aging Study (BLAS) demonstrates Fat Free Mass (FFM) increases up to the 3rd decade of life, after which it remains relatively stable until an accelerated decline, starting at age of 50 for men and 60 for women [1].

Body fat increases until ~ 65 years [2] and redistributes to the abdominal area and visceral organs [3]. The lifetime apex for bone mass (i.e., peak bone mass) occurs in early adulthood [4]. Then, from the 4th to the end of the 8th decade, men and women lose 19% and 39% trabecular BMD, respectively [5]. Collectively body compensation changes contribute to reduced cardiorespiratory capacity [1], and increased risk for cardio-metabolic disease [6,7], sarcopenia [8], osteopenia [9], and frailty [10]. Importantly, physical inactivity exacerbates the age-dependent changes in body composition [11]. As the proportion of people worldwide above 65 yr continues to grow to 16% by 2050 [12], and physical inactivity remains a global problem [11], the prevalence of these conditions will rise, requiring innovative strategies to reduce the socio-economic challenges associated with aging.

Exercise training is recognized as an effective countermeasure to changes in body composition with advancing age [13]; however, optimal exercise training strategies and/or dosing are not fully known. We have previously reported that participants in the Peripheral Remodeling through Intermittent Muscular Exercise (PRIME) trial [14] experienced greater increases in cardiorespiratory and muscular fitness and physical function when a combination of aerobic (AT) and Resistance Training (RT), was preceded by 4 weeks of moderate to high intensity localized muscular exercise (PRIME), compared to moderate to high intensity systemic exercise (AT). The rationale for PRIME was based on: (1) Evidence from the BLAS that accelerated decline in VO₂Peak after 65y is primarily due to reduced peripheral blood flow, skeletal muscle mass and metabolism [1]; and (2) The prevalence of systemic limitations preventing many elderly to achieve an adequate intensity of exercise to stimulate the peripheral barriers [15].

The purpose of the study was: (1) To examine the effects of 12 weeks of exercise training on body composition, including (body weight, Fat and FFM, Bone Mineral Content (BMC), and Bone Mineral Density (BMD)), in elderly at risk for losing functional independence, and (2) To determine if preceding AT/RT, by 4 weeks of PRIME or AT, would contribute to differential body composition responses. It was hypothesized that 12 weeks of exercise training would improve body composition in elderly at risk for losing functional independence, and that the improvements would be greater in those randomized to PRIME compared to AT during the initial 4 weeks of training.

Methods

Recruitment strategy

The goal was to enroll participants over the age of 70, and at risk for losing functional independence based on a peak cardiorespiratory capacity (VO₂peak) of 15-20 mL/kg/min. This range has been shown to be associat-

ed with a threshold for loss of functional independence [16]. The recruitment process involved phone screen interviews to determine participant eligibility. Qualifying participants then attended an orientation visit, including an explanation of study procedures, signing of an informed consent, a review of medical history, and a 6-min walk test [17]. Using previous data from the Louisiana Healthy Aging Study (n = 286, unpublished), subjects who walked between 200 and 450 m in a standard 6-min walk test had a VO₂peak of approximately 15-20 ml/kg/min and could proceed to the baseline testing visits.

Study design

The current report is a 2-arm, prospective pilot randomized trial with participants assigned, to 4 weeks of standard PRIME or AT (Phase I, detailed below). This was followed by 8 weeks of progressive whole-body AT/RT by all participants (Phase II). All individuals gave their informed consent prior to inclusion in the study. The study was conducted at the Pennington Biomedical Research Center, Baton Rouge, LA. The research protocol was reviewed and approved annually by the institutional ethics committee.

Participants

Individuals over the age of 70, and who walked between 200 to 450 m on a 6-min walk test were eligible for participation. Specific exclusion criteria included any unstable or uncontrolled medical conditions. If a volunteer was eligible, they initiated baseline testing (V1) followed by randomization to one of two groups (a) 4 weeks of PRIME or (b) 4 weeks of AT (Phase I). After completion of Phase I, participants completed an intermediate testing assessment (V2), followed by Phase II, which consisted of 8 weeks of a AT/RT based on published physical activity guidelines for older adults [13,18,19]. After phase II, participants completed a final follow-up visit (V3). Both follow-up visits were performed within 5 d of the final Phase I and II training sessions, to minimize any detraining effects. All testing procedures between Phase I and II were completed in less than 7 d.

Exercise interventions

All training sessions lasted 45-60 min, including warm-up, training, and cooldown, and were conducted three times per week for a total of 12 wk. Participants assigned to PRIME, during Phase I, performed eight exercises (see Table 1). Each exercise involved contractions at a moderate load, defined as 40%-50% of maximal voluntary capacity, for a duration of up to 6 min. Participants assigned to AT during Phase I performed whole-body aerobic exercise at 50% of Heart Rate Reserve (HRR) on an Airdyne cycle ergometer (Nautilus, Inc., Vancouver, WA) using both arms and legs for 20 minutes (including a 5-minute warm-up) and then walked on a treadmill for 25 minutes (including a 5-minute cooldown).

Table 1: PRIME exercise protocol.

Exercise	d/wk	Duration (min)	Starting Int	Progression	Cadence	Comments
Calf Raises	3	5	Body Weight	8%-10% of body weight	¼ sec	Both legs
Handgrip	3	5	50% MVC	8%-10% of previous load	¼ sec	Alternating hands
Leg Press	3	6	40-50% MVC	8%-10% of previous load	¼ sec	Both legs
Seated Row	3	5	40-50% MVC	8%-10% of previous load	¼ sec	Both arms
Chest Press	3	5	40-50% MVC	8%-10% of previous load	¼ sec	Both arms
Modified Squat	3	5	Body Weight	8%-10% of previous load	¼ sec	Use of chair or exercise ball
Low Back	3	3	As Tolerated	8%-10% of previous load	¼ sec	Crossed arms
Abdomen	3	3	As Tolerated	8%-10% of previous load	¼ sec	Pads on movement arm on chest

Abbreviations: d/wk = days per week; Int = Intensity; MVC = Maximal Voluntary Contraction

Volume for each exercise was calculated by multiplying weight lifted by the number of repetitions and calculated as volume per exercise, and total volume lifted per session. Heart rate responses to exercise were monitored via Polar heart rate monitors (RS 400; Polar, Kampele, Finland) and data were analyzed for time spent in the prescribed training range and average heart rate achieved during each session. Total energy expenditure was estimated using published metabolic equations [20]. Participants were progressed, as tolerated, over subsequent sessions.

Phase II consisted of 8 weeks of combined AT/RT using established guidelines [13,18,19]. Participants performed 40 minutes of AT (cycle ergometer and treadmill), and one set of 10-15 repetitions using 8 RT exercises. Participants were encouraged to work at 60%-85% of HRR. RT load was increased by 10% for the next session, if the participant could perform the 15th lift of the set. Static stretches were performed after each exercise.

Body composition measurements

Body composition was measured by Dual X-ray Absorptiometry (DXA, QDR 4500A, Hologic Inc., Bedford, MA). Whole body scans were analyzed for Fat Mass, FFM, BMC, and BMD. Whole body scans were regionalized into head, left arm, right arm, left ribs, right ribs, thoracic spine, lumbar spine, pelvis, left leg, and right leg components using established analysis lines and reference points [21].

Additional variables calculated from the raw DXA scores included: (1) Skeletal Muscle Mass Index (SMI), defined by the sum of Appendicular Skeletal Muscle (AMS) divided by height [2,22]; and (2) the sum of the BMC (BMCcore) and BMD scores (BMDcore) calculated from the content and density values of the ribs, thoracic and lumbar spine, and the pelvis.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics Version 24 (IBM, Inc., Armonk, NY). Data are presented as mean (SD) or mean change and 95% confi-

dence intervals, where appropriate. Baseline and demographic data were examined for group differences using t-tests. To address the main hypothesis, the change scores from baseline (V1) to 12 wk (V3) for the primary end points were analyzed. Then, data were analyzed by a repeated measure mixed model, assessing the change after each phase, controlling for the baseline of the outcome under study. Each model assessed group by time interactions, and the main effect.

Results

Participant characteristics

Fifty adults (27 females, 75 ± 4.5 y) were included in this analysis. Baseline data are presented for both PRIME and AT in Table 2. The most common medical diagnoses for participants were: Hypertension (51%), Hypothyroidism (32%), Osteoarthritis (28%), Peripheral Neuropathy (23%), Cancer (16%), and Cardiovascular Disease (15%). The most common medications included anti-hypertensives, and Lipid and thyroid drugs. Medical diagnoses and medications were equally distributed between groups.

Training volume

Bi-weekly exercise data are presented in Table 3. The data include average intensity during each session, progression across the study for volume lifted, and estimated energy expenditure (Phase I AT only). The intensity of training for PRIME during Phase I ranged from 41.2 ± 21.5% to 45.8 ± 24.3% HRR compared to 56.7 ± 15.5% to 64.5 ± 15.5% HRR for AT (Group comparison, p = 0.01).

Body composition

Body weight did not change. Body fat % decreased in both groups (p = 0.005), and to a greater extent in AT (Fat %: PRIME: Δ -0.47 [95% CI: -0.98 to 0.04]; AT: Δ -0.98% [95% CI: -1.44 to -0.50], p = 0.009). The reduction in fat mass was greater in AT (Fat Mass: PRIME: Δ -483 g [95% CI: -748 to 252]; AT: Δ -1039 g [95% CI: -1546 to -666], p = 0.05). Lean mass increased in both groups (p = 0.03), (Lean Mass: PRIME: Δ 327 g [95% CI: -166 to 820];

Table 2: Study Participants Baseline (V1) Characteristics.

	All (n = 50)	PRIME (n = 23)	AT (n = 27)
Age	75.0 ± 4.52	75.0 ± 3.98	75.0 ± 4.97
Gender			
Female	33	16	17
Resting Hemodynamics			
Heart Rate	67.5 ± 8.21	67.0 ± 7.02	68.5 ± 9.31
Systolic Blood Pressure	130.8 ± 13.23	131.9 ± 13.95	130.3 ± 12.75
Diastolic Blood Pressure	72.8 ± 8.83	73.6 ± 10.16	72.1 ± 7.53
Body Composition			
Height	164.58 ± 8.54	165.2 ± 9.06	163.7 ± 8.18
Weight	80.97 ± 16.45	82.48 ± 20.17	79.7 ± 12.48
Fat Mass	31.26 ± 8.77	29.98 ± 7.11	32.36 ± 9.36
Lean Mass	50.43 ± 11.46	49.14 ± 10.95	51.73 ± 11.64
SMI (AMS/m ²)	6.44 ± 1.59	6.25 ± 1.49	6.60 ± 1.67
Bone Mineral Density			
Arms	0.76 ± 0.11	0.73 ± 0.11	0.77 ± 0.11
Ribs	0.68 ± 0.12	0.67 ± 0.10	0.70 ± 0.14
Thoracic Spine	0.97 ± 0.16	0.94 ± 0.17	1.00 ± 0.15
Lumbar Spine	1.04 ± 0.22	1.02 ± 0.24	1.06 ± 0.21
Pelvis	1.20 ± 0.17	1.18 ± 0.17	1.22 ± 0.17
Legs	1.20 ± 0.19	1.18 ± 0.20	1.22 ± 0.18

Values are means ± SD

Table 3: Bi-weekly exercise training data.

	Phase 1		Phase 2			
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
PRIME	Weeks 1&2	Weeks 3&4	Weeks 5&6	Weeks 7&8	Weeks 9&10	Weeks 11&12
Volume Lifted, lbs	37291 ± 7637	42143 ± 7603	10965 ± 2917	12837 ± 3024	13172 ± 3288	14305 ± 2699
Intensity, % HRR	41.2 ± 21.5*	45.8 ± 24.3*	60.0 ± 19.2	65.6 ± 17.3	70.3 ± 16.7	71.0 ± 19.3
EE, Kcal	N/A	N/A	130 ± 33.7	162 ± 41.0	177 ± 45.5	188 ± 47.0
AT	Weeks 1&2	Weeks 3&4	Weeks 5&6	Weeks 7&8	Weeks 9&10	Weeks 11&12
Volume Lifted, lbs	N/A	N/A	11084 ± 3260	12867 ± 3642	14304 ± 4061	15080 ± 4412
Intensity, % HRR	56.7 ± 15.5	64.5 ± 15.5	62.0 ± 19.5	65.9 ± 16.2	68.3 ± 14.5	69.1 ± 15.3
EE, Kcal	160 ± 36.5	191 ± 39.9	166 ± 35.0	183 ± 38.5	196 ± 48.1	206 ± 52.5

Abbreviations: SD = Standard Deviation, HRR = Heart Rate Reserve, EE = Energy expenditure. N/A = Not Applicable.

Data presented bi-weekly with per-session (3 sessions per week) averages. *p < 0.05, Phase I group comparison for exercise intensity

AT: Δ 232 g [95% CI: -350.88 to 815.24]). Finally, SMI increased in both groups (p = 0.05) from V1 to V3.

Bone mineral content and bone mineral density

Data for BMC and BMD are presented in [Table 4](#) and [Table 5](#). The Tables provide p-values for time effect, group by time interaction, and group differences for the change scores from V2 to V1, and from V3 to V1. For those in PRIME, BMC improved at the ribs, thoracic spine, and pelvis. For AT BMC decreased at the ribs, and Thoracic Spine. Change scores (V3-V1) between groups were: (1) **Arms** (PRIME: -2.81 g [95% CI: -5.63 to 0.01]; AT: -2.02 g [95% CI: -4.62 to 0.59], p = 0.68); (2)

Ribs (PRIME: 5.61 g [95% CI: 0.73 to 10.49]; AT: -4.41 g [95% CI: -8.92 to 0.09], p = 0.004); (3) **Thoracic Spine** (PRIME: 6.31 g [95% CI: -3.11 to 15.74]; AT: -8.92 g [95% CI: -17.62 to -0.23], p = 0.02), **Lumbar spine** (PRIME: 2.64 g [95% CI: -1.68 to 6.96]; AT: 5.83 g [95% CI: 1.84 to 9.82], p = 0.28), **Pelvis** (PRIME: 1.72 g [95% CI: -3.68 to 7.13]; AT: -1.15 g [95% CI: -6.13 to 3.84], p = 0.44), **Legs** (PRIME: 3.46 g [95% CI: -4.58 to 11.49]; AT: -2.67 g [95% CI: -10.09 to 4.74], p = 0.27), and for **BMCcore** (PRIME: 14.54 g [95% CI: 4.48 to 24.61]; AT: -8.28 g [95% CI: -17.57 to 1.01], p = 0.002).

BMD improved at the ribs, thoracic and lumbar spine, and pelvis, for PRIME, and AT at the pel-

Table 4: Bone mineral content (grams) for V1, V2, and V3.

	Group	Bound	AT			PRIME			p-values			
			V1	V2	V3	V1	V2	V3	Time Effect	Grp* Time	Grp Diff for Δ V2-V1	Grp Diff for Δ V3-V1
Arms	Mean		167.40a	166.33	165.51	167.40a	166.74	90.58	0.89	0.76	0.84	0.68
	95% CI	LB	167.40	163.25	162.90	167.40	163.41	170.08				
		UB	167.40	169.41	168.20	167.40	161.61	167.27				
Ribs	Mean		86.35a	86.66	82.49	86.35a	89.94	91.31	0.03	0.02	0.18	0.004
	95% CI	LB	86.35	82.24	78	86.35	85.15	86.45				
		UB	86.35	91.09	86.97	86.35	94.74	96.17				
Thoracic	Mean		136.94a	138.96	127.53	136.94a	141.3	144.03	0.03	0.01	0.52	0.02
	95% CI	LB	136.94	130.81	117.9	136.94	132.45	133.38				
		UB	136.94	147.12	137.15	136.94	150.14	154.26				
Lumbar	Mean		55.72a	58.32	62.01	55.72a	57.37	58.34	0.13	0.42	0.64	0.28
	95% CI	LB	55.72	53.82	56.67	55.72	52.42	51.17				
		UB	55.72	62.82	67.36	55.72	62.17	62.75				
Pelvis	Mean		225.62a	229.36	224.71	225.62a	228.28	229.86	0.26	0.05	0.02	0.44
	95% CI	LB	225.62	215.61	220.14	225.62	224.12	218.33				
		UB	225.62	225.12	229.28	225.62	224.90	234.82				
Legs	Mean		451.56a	452.39	448.68	451.56a	458.71	455.07	0.80	0.50	0.36	0.27
	95% CI	LB	451.56	443.55	441.34	451.56	449.12	446.95				
		UB	451.56	461.24	456.34	451.56	510.77	463.19				
BMCcore	Mean		505.23a	503.15	496.06	505.23a	518.79	520.82	0.16	0.001	0.02	0.002
	95% CI	LB	505.23	494.45	480.38	505.23	509.35	510.95				
		UB	505.23	511.84	505.16	505.23	528.22	530.70				


Abbreviations: Grp = Group; Diff = Difference; CI=Confidence Interval; V1 = Visit 1; V1 = Visit 1; V2 = Visit 2; V3 = Visit 3; Δ = Change score; LB = Lower Bound; UB = Upper Bound; BMC = Bone Mineral Content.

Table 5: Bone mineral density (g/cm³) for V1, V2, and V3.

	Group	Bound	AT			PRIME			p-values			
			V1	V2	V3	V1	V2	V3	Time Effect	Grp*Time	Grp Diff for Δ V2-V1	Grp Diff for Δ V3-V1
Arms	Mean		0.76a	0.74	0.75	0.76a	0.75	0.75	0.12	0.40	0.19	0.74
	95% CI	LB	0.76	0.73	0.73	0.76	0.74	0.73				
		UB	0.76	0.75	0.76	0.76	0.77	0.76				
Ribs	Mean		0.68a	0.69	0.68	0.68a	0.70	0.71	0.005	0.06	0.28	0.05
	95% CI	LB	0.68	0.68	0.66	0.68	0.69	0.69				
		UB	0.68	0.71	0.70	0.68	0.72	0.73				
Thoracic	Mean		0.97a	1.00	0.98	0.97a	1.00	1.04	0.11	0.003	0.62	0.007
	95% CI	LB	0.97	0.97	0.95	0.97	0.96	1.01				
		UB	0.97	1.03	1.01	0.97	1.03	1.08				
Lumbar	Mean		1.05a	1.05	1.05	1.05a	1.09	1.09	0.69	0.03	0.18	0.05
	95% CI	LB	1.05	1.03	1.02	1.05	1.06	1.06				
		UB	1.05	1.08	1.08	1.05	1.12	1.12				
Pelvis	Mean		1.12a	1.20	1.21	1.12a	1.20	1.22	0.04	0.73	0.62	0.38
	95% CI	LB	1.12	1.18	1.20	1.12	1.18	1.21				
		UB	1.12	1.21	1.23	1.12	1.22	1.25				
Legs	Mean		1.20a	1.19	1.18	1.20a	1.20	1.20	0.60	0.33	0.49	0.31
	95% CI	LB	1.20	1.18	1.17	1.20	1.18	1.18				
		UB	1.20	1.20	1.21	1.20	1.22	1.21				

BMDcore	Mean		3.90a	3.93	3.92	3.90a	3.99	4.07	0.23	0.0001	0.20	0.001
	95% CI	LB	3.90	3.88	3.87	3.90	3.93	4.04				
		UB	3.90	3.98	3.97	3.90	4.01	4.13				

Abbreviations: Grp = Group; Diff = Difference; CI=Confidence Interval; V1 = Visit 1; V1 = Visit 1; V2 = Visit 2; V3 = Visit 3; Δ = Change score; LB = Lower Bound; UB = Upper Bound; BMD = Bone Mineral Density



vis. Change scores (V3-V1) between groups were:(1) **Arms**: PRIME: 0.030 g/cm³ [0.070 to 0.054]; AT: -0.004 g/cm³ [-8.69-2.15], p = 0.04, (2) **Ribs** (PRIME: 0.036 g/cm³ [95% CI: 0.014 to 0.057]; AT: -0.003 g [95% CI: -0.022 to 0.016], p = 0.009); (3) **Thoracic Spine** (PRIME: 0.071 g/cm³ [95% CI: 0.039 to 0.103]; AT: 0.009 g/cm³ [95% CI: -0.018 to 0.037], p = 0.0001), **Lumbar spine** (PRIME: 0.046 g/cm³ [95% CI: 0.013 to 0.079]; AT: 0.002 g/cm³ [95% CI: -0.027 to 0.030], p = 0.82), **Pelvis** (PRIME: 0.025 g/cm³ [95% CI: 0.006 to 0.043]; AT: 0.014 g/cm³ [95% CI: -0.003 to 0.030], p = 0.24), **Legs** (PRIME: -0.002 g/cm³ [95% CI: -0.017 to 0.014]; AT: -0.018 g/cm³ [95% CI: -0.031 to -0.004], p = 0.23), and for **BMDcore** (PRIME: 0.206 g/cm³ [95% CI: 0.140 to 0.271]; AT: 0.001 g/cm³ [95% CI: -0.058 to 0.058], p = 0.0001).

Individual changes in **BMCcore** and **BMDcore** are depicted in [Figure 1a](#) and [Figure 1b](#). Those randomized to PRIME had a greater likelihood of improvement in **BMCcore** and **BMDcore** after Phase II, with 70% of PRIME versus 25% of AT ([Figure 1a](#)) and 78% of PRIME versus 55% of AT ([Figure 1b](#)) demonstrating improvements, respectively.

Discussion

The unique contributions of this study are 2-fold: (1) Participants whom initiated training with PRIME experienced greater improvements in BMC and BMD, and (2) Starting an exercise program for older adults at risk of losing independence, using a low intensity systemic, yet moderate to high intensity localized exercise stimulus, represent a novel approach which may be beneficial to those with cardiorespiratory or other systemic limitations.

Body composition

Both groups decreased percent body fat. The ~1% decrease in percent fat is similar to those observed in the original study from Pollock's laboratory, comparing AT and RT responses in 70 to 79-year-olds [23]. Similarly changes in fat mass were greater in those who performed AT only [23]. Importantly, both groups increased SMI. These changes are relevant in that SMI is used to identify those considered sarcopenic [22,24]. The PRIME study was not powered to address this issue directly, so we remain conservative, only to suggest exercise training serves as a countermeasure to advancing sarco-




Figure 1b: Individual **BMDcore** changes after 12 weeks of exercise training.

penia. Future studies should address the value of SMI changes in regard to maintaining independence. The small changes in FFM may be due to neural adaptations, rather than hypertrophy [25].

Bone mineral content and density

A unique finding within this report is the observed increases in BMC and BMD, particularly in the PRIME group. The findings BMC and BMD increased to a greater extent in PRIME versus AT suggests the differential loading conditions in Phase I were important in the overall training response. The confidence in the observed findings may be explained as follows: (1) The number of targeted exercises used, in PRIME, during PHASE I; (2) The progression in the mechanical loading conditions within Phase I, in the targeted areas; (3) The similarities in the responses between PRIME and AT in the pelvic area; and (4) The individual responses in BMC and BMD in those randomized to PRIME or AT. In regard to the first point, six of 8 PRIME exercises aimed to target core strength, involving many of the areas and sites measured with the DXA. The delivery of these exercises was such that the load and cycle number were high, as shown to be required [26]. Moreover, the exercises were developed to ensure sufficient overlap on major muscle groups and joints. Thus, most muscle groups, in PRIME, received multiple loading conditions, within the same session, potentially optimizing the stimulus for adaptation. In addition, those involved in PRIME had an increase in volume/load of ~14% from week 1 & 2 to week 3 & 4,

in Phase I. Consequently, the specific sites received considerable and progressive stimulation. Arguably, the AT group did not receive such a consistent stimulus, during Phase I, except for the Pelvic area, due to its involvement in cycling and walking. Finally, PRIME participants responded to the training period in a more favorable manner, with 16 and 18 out of 23 PRIME participants reporting an increase in **BMCcore** and **BMDcore** vs. 7 and 15 out of 27 of AT participants, respectively.

The observed increases in this study, seen within 3 months, are of similar magnitude to studies with training programs over 6 months. For example, a recent systematic review examining BMD changes in older adults following exercise training showed the earliest evidence for change around 4 months [26]. In fact, the first study to report a change in BMC of ~5% after 6 months of walking at an intensity of 70-90% VO_2max was performed in post-menopausal women [27]. However, most studies that have examined the effects of AT on BMC and BMD reported no increase unless high mechanical loads are generated at the bone [26]. RT provides more consistent evidence of increased BMC and BMD in older populations, after 6 months. The gains typically involve only the areas that receive the mechanical load [26].

The greater improvements in BMC and BMD, in PRIME, evident after 12 weeks are, intriguing. During Phase II of the intervention both groups trained at similar intensities and progressed equally (Table 3): We can only speculate the initial training period provided a su-

perior signal for adaptation that subsequently carried through phase II. It is premature to suggest underlying mechanisms for the changes, but evidence indicates the skeletal muscle secretome is sensitive to mechanical contractions [28]. The subsequent effects of muscle contractions may produce specific myokines involved in triggering a number of pathways contributing to end-organ responses, including bone [29].

Clinical relevance

According to Wolfe: “Elderly individuals particularly women, are often too weak to perform the intensity of exercise necessary to induce the same magnitude of physiologic adaptations that occur in younger” [30]. In fact, many individuals are not able to perform at a high enough intensity of exercise to induce adaptations because of systemic limitations. The current study offers a novel approach that allows for significant changes in physiologic and functional phenotypes, if training is initiated with a low-mass, high-repetition/volume mechanical loading scheme under a relatively low systemic strain, and followed by training guidelines as recommended by American College of Sports Medicine/American Heart Association guidelines) [19].

Study limitations

Exercise guidelines for older individuals include recommendations for a well-rounded training program applied in the context of the participant’s needs, goals, and baseline functional abilities, and with a gradual progression in duration and intensity of effort for cardio-respiratory, muscular strength and endurance, and flexibility components [13,18,19]. The recommendations provide little insight how to incorporate principles of specificity, overload and progression into the regimen in order to optimize functional benefits. Implementation usually starts with a moderate intensity systemic exercise stimulus, with a gradual increase in time and intensity as tolerated and inclusion of RT later in the program. This study is thus delimited in our interpretation of the order which is common practice, i.e. AT first, followed by gradual increase of other training components. We do not know how individuals would compare if PRIME were to be matched with RT (or any other order of presentation).

The selection of 4 weeks of PRIME is based on previous studies which focused on the effects of a local training stimulus on vasodilatory function [31,32]. PRIME has not been extended beyond 4 weeks, to determine if the responses differ. However, within that argument it is perhaps key to focus on the consistent improvements observed in BMC and BMD, with PRIME, in this vulnerable population. Not only are the improvements in functional ability, relevant, as reported, but the present study could potentially support the use of PRIME as an early intervention for those who are osteopenic.

Conclusion

Exercise training for older adults at risk of losing independence results in beneficial changes in body composition. Importantly, this study highlights the feasibility and superiority of initiating exercise training using a low intensity systemic, yet moderate to high intensity localized exercise stimulus, as it relates to bone health. As such these data support further research to examine the need for a shift in the paradigm of initiating exercise training in the elderly from the traditional moderate to high intensity systemic exercise stimulus, toward the use of moderate to high intensity localized exercise.

Competing Interests

The authors declare that they have no competing interests.

Authors’ Contributions

MAW and JDA conceived and designed the study, analyzed data, interpreted the findings and contributed to writing the manuscript; MCS executed the exercise programs, analyzed data, interpreted the findings, and contributed to writing the manuscript; NMJ and DPC helped in the design of the study, interpreted the findings and contributed to writing the manuscript. CPE, TSC, ER, WEK, assisted in interpreting the findings and contributed to writing the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of the presentation of authors.

This work was supported by the National Institutes of Aging (grant no. 1RC1AG035822, 2009).

References

1. Fleg JL, Morrell CH, Bos AG, Brant LJ, Talbot LA, et al. (2005) Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation* 2: 674-682.
2. Wilson PW, Kannel WB (2002) Obesity, diabetes, and risk of cardiovascular disease in the elderly. *Am J Geriatr Cardiol* 11: 119-123.
3. Hunter GR, Gower BA, Kane BL (2010) Age Related Shift in Visceral Fat. *Int J Body Compos Res* 8: 103-108.
4. Baxter-Jones AD, Faulkner RA, Forwood MR, Mirwald RL, Bailey DA (2011) Bone mineral accrual from 8 to 30 years of age: an estimation of peak bone mass. *J Bone Miner Res* 26: 1729-1739.
5. Wilsgaard T, Emaus N, Ahmed LA, Grimnes G, Joakimsen RM, et al. (2009) Lifestyle impact on lifetime bone loss in women and men: the Tromsø Study. *Am J Epidemiol* 169: 877-886.
6. Eckel RH, Grundy SM, Zimmet PZ (2005) The metabolic syndrome. *Lancet* 365: 1415-1428.
7. Frayn KN (2000) Visceral fat and insulin resistance—causative or correlative? *Br J Nutr* 83: 71-77.
8. Evans WJ, Campbell WW (1993) Sarcopenia and age-related changes in body composition and functional capacity. *J Nutr* 123: 465-468.
9. Nguyen TV, Kelly PJ, Sambrook PN, Gilbert C, Pocock NA,

- et al. (1994) Lifestyle factors and bone density in the elderly: implications for osteoporosis prevention. *J Bone Miner Res* 9: 1339-1346.
10. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, et al. (2001) Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 56: 146-156.
11. Booth FW, Roberts CK, Thyfault JP, Rueggsegger GN, Toedebusch RG (2017) Role of Inactivity in Chronic Diseases: Evolutionary Insight and Pathophysiological Mechanisms. *Physiol Rev* 97: 1351-1402.
12. National Institutes of Health, Organization WH. Global Health and Aging.
13. American College of Sports Medicine, Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, Minson CT, et al. (2009) American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc* 41: 1510-1530.
14. Allen JD, Vanbruggen MD, Johannsen NM, Robbins JL, Credeur DP, et al. (2018) PRIME: A Novel Low-Mass, High-Repetition Approach to Improve Function in Older Adults. *Med Sci Sports Exerc* 50: 1005-1014.
15. Paterson DH, Warburton DE (2010) Physical activity and functional limitations in older adults: a systematic review related to Canada's Physical Activity Guidelines. *Int J Behav Nutr Phys Act* 7: 38.
16. Cress ME, Meyer M (2003) Maximal voluntary and functional performance levels needed for independence in adults aged 65 to 97 years. *Phys Ther* 83: 37-48.
17. Allen JD, Robbins JL, VanBruggen MD, Credeur DP, Johannsen NM, et al. (2013) Unlocking the barriers to improved functional capacity in the elderly: rationale and design for the "Fit for Life trial." *Contemp Clin Trials* 36: 266-275.
18. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, et al. (2011) Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 43: 1334-1359.
19. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, et al. (2007) Physical activity and public health in older adults. Recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation* 116: 1094-1105.
20. Riebe D, Ehrman JK, Liguori G, Magal M, American College of Sports Medicine (2017) ACSM Guidelines for Exercise Testing and Prescription. (10th edn), Philadelphia: Wolters Kluwer, 192-193.
21. Lohman M, Tallroth K, Kettunen JA, Marttinen MT (2009) Reproducibility of dual-energy x-ray absorptiometry total and regional body composition measurements using different scanning positions and definitions of regions. *Metabolism* 58: 1663-1668.
22. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, et al. (2010) Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 39: 412-423.
23. Hersey WC, Graves JE, Pollock ML, Gingerich R, Shireman RB, et al. (1994) Endurance exercise training improves body composition and plasma insulin responses in 70- to 79-year-old men and women. *Metabolism* 43: 847-854.
24. Locquet M, Beaudart C, Bruyère O, Kanis JA, Delandshere L, et al. (2018) Bone health assessment in older people with or without muscle health impairment. *Osteoporos Int* 29: 1057-1067.
25. Moritani T, deVries HA (1980) Potential for gross muscle hypertrophy in older men. *J Gerontol* 35: 672-682.
26. Gómez-Cabello A, Ara I, González-Agüero A, Casajús JA, Vicente-Rodríguez G (2012) Effects of training on bone mass in older adults: a systematic review. *Sports Med* 42: 301-325.
27. Dalsky GP, Stocke KS, Ehsani AA, Slatopolsky E, Lee WC, et al. (1988) Weight-bearing exercise training and lumbar bone mineral content in postmenopausal women. *Ann Intern Med* 108: 824-828.
28. Hamrick MW (2012) The skeletal muscle secretome: an emerging player in muscle-bone crosstalk. *Bonekey Rep* 1: 60.
29. Wolfe RR (2006) The underappreciated role of muscle in health and disease. *Am J Clin Nutr* 84: 475-482.
30. Pedersen BK (2013) Muscle as a secretory organ. *Compr Physiol* 3: 1337-1362.
31. Allen JD, Geaghan JP, Greenway F, Welsch MA (2003) Time-Course of Vascular Adaptations following Localized Short-Term Exercise Training. *Med Sci Sports Exerc* 35: 847-853.
32. Dobrosielski DA, Greenway F, Welsh D, Jazwinski SM, Welsch MA, et al. (2009) Modification of vascular function in elderly men following handgrip exercise training. *Med Sci Sports Exerc* 41: 1429-1435.