



RESEARCH ARTICLE

Laparoscopic Choledochal Cyst Excision and Roux-En-Y Hepaticojejunostomy in Children: A Single-Institution Prospective Cohort Study

Nguyen Thanh Xuan¹, Ho Huu Thien¹, Phan Hai Thanh¹, Pham Anh Vu², Dang Nhu Thanh¹, Nguyen Du Vinh¹, Nguyen Huu Son^{3*} and Pham Nhu Hiep¹

¹Department of Pediatric and Abdominal Emergency Surgery, Hue Central Hospital, Vietnam

²Department of Surgery, Hue University of Medicine and Pharmacy, Vietnam

³Pediatric Center, Hue Central Hospital, Vietnam



*Corresponding author: Nguyen Huu Son, Pediatric Center, Hue Central Hospital, 16 Le Loi street, Hue city, Vietnam

Abstract

Purpose: Laparoscopic cyst excision and Roux-en-Y hepaticojejunostomy is gaining popularity as a treatment for choledochal cyst in children. The aim of this study is to determine the feasibility and safety of the laparoscopic excision with Roux-en-Y hepaticojejunostomy, and evaluate the short-term outcomes after treatment for children with choledochal cyst.

Methods: A prospectively of 51 consecutive pediatric patients undergoing laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy performed by one surgical team at Hue Central Hospital from June 2012 to December 2017.

Results: The mean operative time was 214.7 ± 67.95 minutes (range, 100~360 minutes), including the time for intraoperative cholangiography. There were two children requiring blood transfusion. Time to first flatus was 40.35 ± 28.55 hours in average. The mean time to drain removal was 2.89 ± 1.02 days. Mean postoperative hospital stay was 9.31 ± 3.43 days. 6/51 cases have early complications, including 2 cases of pancreatitis and 4 cases of bile leakage. Most of cases (90.5%) were classified as good after 10 days to 3 months of follow-up.

Conclusion: Laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy was feasible and safe in children. The short-term outcomes were good in most cases.

Keywords

Laparoscopic excision, Roux-en-Y hepaticojejunostomy, Choledochal cyst, Children

Introduction

Choledochal cysts are a rare congenital cystic dilation of the biliary tract, first described by Vater and Ezler in 1723. They present primarily in female infants and young children and are more prevalent in East Asian populations. Although benign, choledochal cysts can be associated with serious complications including malignant transformation, cholangitis, pancreatitis, and cholelithiasis [1,2]. Hence, complete surgical excision of the choledochal cyst and enterobiliary reconstruction has been recommended as the standard treatment.

Choledochal cyst excision and biliary enteric reconstruction constitutes the best therapy for choledochal cyst [3]. Comparing with open procedure, laparoscopic procedure has been proven to have shorter hospital stay and lower morbidity of anastomotic stenosis, bile leakage, intrahepatic stone formation, cholangitis, pancreatic leak, intestinal obstruction, and re-operation [4,5]. Thus, as a safe, efficacious, and minimally invasive procedure, laparoscopic cyst excision and Roux-en-Y hepaticojejunostomy has become a common procedure for pediatric choledochal cyst in many medical centers [6]. With increased familiarity with the laparoscopic anatomy of the biliary tract and advances in minimally invasive techniques, surgeons have ventured further to operate on technically difficult cases such as choledochal cyst that were until recently managed by laparot-

omy [7]. Our preliminary result shows that laparoscopic excision of choledochal cyst and Roux-en-y hepaticojejunostomy is effectiveness in pepdiatric patients [8]. The purpose of this study was to determine the feasible and safe of the laparoscopic excision with Roux-en-Y hepaticojejunostomy, and evaluate the short-term outcomes after treatment for children with choledochal cyst.

Materials and Methods

Patients

A prospective study involving 51 pediatric patients diagnosed with choledochal cysts at Hue Central Hospital from June 2012 to December 2017. All patients underwent laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy. All procedures were performed by the same surgery team. This study was approved by the ethics committee review board of our hospital.

The variables investigated were demographic information of all patients, operative details and outcomes such as operation time, intraoperative blood transfusion, postoperative time to first flatus and postoperative hospital stay, and postoperative complications.

Operative technique

The procedure of laparoscopic choledochal cyst exci-



Figure 1: Transection of cystic duct.




Figure 2: Distal common bile duct was transected between two hemoclips.

sion is summarized below:

1) Under general anesthesia, the patient was intubated and placed in reverse Trendelenburg position.

2) Four port procedure was used as the trocars were located at middle of the supra-umbilicus (10 mm), right hypochondrium (2 ports 5 mm), right side of the abdomen, and left hypochondrium (5 mm), respectively.

3) A monopolar electrocautery hook was used to dissect the choledochal cyst and the gallbladder. The cystic duct and artery were transected and the gallbladder was separated from the cyst but was not removed from the liver bed (Figure 1). The gallbladder was sutured with 2/0 silk from the fundus to the right diaphragmatic area to achieve liver retraction. The cyst was dissected down to the distal tapered end of the common bile duct, and it was then transected between 2 hemoclips (Figure 2). The upper part of the cyst was further dissected up to the common hepatic duct. When severe adhesion around the cyst was encountered, thunderbeat (Olympus) was used for dissecting.

4) The jejunum was transected with an endoscopic stapler 25 cm away from the ligament of Treitz for hepaticojejunostomy. A side-to-side enteroenterostomy was created by the exteriorization of the small bowel via the enlarged umbilical port between the loop 40 cm distal from the Roux loop and the afferent loop from the Treitz ligament. The distal end was advanced in a retrocolic manner, and the Roux loop was brought to the liver hilum. The cyst was transected and extracted out of the umbilical site with the gallbladder (Figure 3). When the bile duct was opened, it was observed that the biliary bifurcation has been reached (Figure 3). The anastomosis was created following an enterotomy with interrupted Vicryl 4/0 sutures) (Figure 4).

5) A redon suction drain was placed posterior to the hepaticojejunostomy after completion of cholecystectomy which was remove on 3rd day or bile drainage less than 3 ml/h.

6) When two bile ducts (left and right hepatic ducts)




Figure 3: Common hepatic duct was transected at the hilum.




Figure 4: Hepato-jejuno anastomosis with interrupted suture.

Table 1: Patient characteristics and laboratory tests.

Variables	All patients (n = 51)
Gender	
Female	41 (80.4%)
Male	10 (19.6%)
Age (mean, range)	3.55 years (2.5 months ~15 years)
Symptoms	
Abdominal pain	38 (74.5%)
Jaundice	17 (33.3%)
Fever	18 (35.3%)
Nausea, vomiting	25 (49.0%)
Abdominal mass	3 (5.9%)
Laboratory findings	
Increased serum AST/ALT	25 (49.0%)
Increased total bilirubin	45 (88.2%)
Leukocytosis	17 (33.3%)
Operative time (mins, range)	214.7 ± 67.95 (100 ~ 360)

Table 2: Surgical outcomes of laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy.

Variables	All patients (n = 51)
Number of patient requiring blood transfusion	2 (3.9%)
Blood transfusion volume (ml)	125.0 ± 35.36
Time to first flatus (hours)	40.35 ± 28.55
Time to drain removal (days)	2.89 ± 1.02
Hospital stay (days)	9.31 ± 3.43
Complications	
Biliary leakage	4
Fluid collection	0
Pancreatitis	2
Outcome classification (Terblanch)	n = 42 (9 patients were lost to follow-up)
I	38 (90.5%)
II	3 (7.1%)
III	0
IV	1 (2.4%)

were encountered in the Todani-IVa type cases, a ductoplasty was performed and followed by choledochostomy.

Postoperative management and follow-up of patients

Patients usually started a liquid diet on postoperative day 1. Jackson-Pratt drains were removed if ultrasound that were performed on postoperative day 4 showed no evidence of abnormal fluid collection. Patients revisited the outpatient department 2 weeks after discharge and were followed up at 3 months to monitor postoperative complications. For follow-up, a routine complete blood count, liver function test, and ultrasound or CT were performed to evaluate complications, such as pancreatitis, cholangitis, choledocholithiasis, or anastomosis stenosis.

Clinical outcome was determined according to the Terblanche classification [9]: Grade I, no biliary symptoms; grade II, transitory symptoms and no current symptoms; grade III, biliary symptoms requiring medical therapy; and grade IV, recurrent biliary symptoms requiring correction or related to death. Terblanche class IV constituted a poor result. Terblanche I, II, and III constituted a clinical success with excellent, good, and fair results, respectively.

Data analysis

For a descriptive analysis, the frequency or the mean and standard deviation were calculated for each variable. For other continuous variables, independent sample t-tests were applied to compare the data from the children and adult groups. Their respective p-values and corresponding confidence intervals were provided by SPSS Version 18.0 (SPSS Inc., Chicago, Illinois, USA). The statistical significance was set at $p < 0.05$.

Results

Forty-one patients were females and 10 were males. The mean age at the time of surgery was 3.55 years (range, 2.5 months~15 years). The patient characteristics and the operative features were seen in Table 1. Abdominal pain was noted in 74.5%, nausea and vomiting in 25 cases (49%), jaundice in 17 cases (33.3%), high serum hepatic enzyme levels in 25 children (49.0%). The increase in serum bilirubin level along with different symptoms of jaundice was seen in 45 children (88.2%).

The mean operative time was 214.7 ± 67.95 minutes (range, 100~360 minutes), including the time for intraoperative cholangiography. The conversion to open did not happen in any cases.

Table 2 shows the surgical outcomes of laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy. There were two children requiring blood transfusion with the total volume of packed red blood cells of 125.0 ± 35.36 ml. Time to first flatus was 40.35 ± 28.55

hours in average. The mean time to drain removal was 2.89 ± 1.02 days. Mean postoperative hospital stay was 9.31 ± 3.43 days. 6/51 cases having early complications, including 2 cases of pancreatitis and 4 cases of bile leakage. Conservative treatment was initially implemented which resulted in complete resolution in 5 cases. Reoperative was finally required in 1 cases with persistent bile leakage. Most of cases were classified as good after hospital discharge to 3 months of follow-up.

Discussion

Choledochal cyst is a rare congenital anomaly of the biliary system that commonly affects the Asian population [10]. The estimated incidence of choledochal cyst and anomalous pancreatobiliary ductal union is 0.3% and 4.1%, respectively, according to a large, multicenter study [11]. The clinical presentation of choledochal cyst differs from children to adults. In adults, this disease entity can manifest cholangitis, pancreatitis, choledocholithiasis, or biliary malignancy, such as cholangiocarcinoma or gallbladder cancer [11]. Complete excision of the cyst is mandatory because of the risk of developing complications or malignancies.

With growing interest in cosmetics, laparoscopic approaches in hepatobiliary surgery have become inevitable. Laparoscopic hepaticojejunostomy anastomosis still remains one of the most sophisticated and challenging surgeries to date [12]. Since the first report of laparoscopic choledochal cyst excision in 1995 [13], studies regarding this approach have reported that laparoscopic surgery is safe and feasible in the treatment of choledochal cysts.

Choledochal cyst is characterized by its remarkably higher incidence in the female patients together with the regional preference. In this study also, we found more cases of female patients, the female/male ratio in our study was 4.0/1. This was similar to other studies. Sheng Q (2017) showed a female/male ratio of 15/3 = 5 [14]. In the study of Liu Y, et al., the female/male ratio was 29/6 = 4.83 [5]. Since choledochal cyst is more common in female, the aesthetic advantages of laparoscopic procedure are more relevant. Patient's age ranged from 2.5 months to 15 years (Table 1). The mean age was 3.55 years. The results of our study were similar to other studies. Huang CS, et al. (2010) reported 101 cases of choledochal cysts in which 42 cases were children (41.6%) and 59 cases were adults (58.4%). The mean age was 8.9 years 3.7 years for children [15].

We performed HJ with Rous-en-Y technique in all cases. Operative time ranged from 100 to 320 mins. Mean operative time was 214.7 ± 67.95 mins. Minimizing operative time and maintaining outcomes at the same time has always been a challenge to surgeons, especially in young children and infants. If surgery is too lengthy, comorbidity rate will increase. To shorten operative time, multiple factors are required including competent skills of surgeon, assistants and nurses,

close coordination with anesthesiologists and adapted instruments. Therefore, operative time in the beginning of the learning curve of laparoscopic procedure is longer. Operative time of laparoscopic procedure was also longer compared to open procedure in recent studies. Our operative time was comparable to the study of Liem NT (2012) with 400 patients, the operative time in hepatoduodenal anastomosis group was 164.8 mins, and in hepatojejunal anastomosis group was 220 mins [16]. According to Liu Y, et al., mean operative time of laparoscopic procedure was 249 mins, longer than open procedure with mean operative time of 132 mins [5].

One of the complications of laparoscopic choledochal cyst excision is bleeding requiring intraoperative blood transfusion. Liem NT (2012) reported four cases with bleeding requiring intraoperative blood transfusion, three cases with transection of both hepatic ducts and one case with right hepatic duct perforation [16]. Tang ST, et al. noted 8 cases requiring intraoperative blood transfusion [17].

Regarding the early postoperative complications, we noted 6/51 cases having early complications, including 2 cases of pancreatitis and 4 cases of bile leakage. Conservative treatment was initially implemented which resulted in complete resolution in 5 cases. Reoperative was finally required in 1 cases with persistent bile leakage. Reoperative was finally required in 1 cases with persistent bile leakage. Ono, et al. [18] reported liver dysfunction, dilatation of intrahepatic bile ducts, recurrent abdominal pain, and biliary tract malignancy as long-term complications after choledochal cyst excision and hepaticojejunostomy. The laparoscopic approach can magnify the operative view, which may assist in dissecting the whole choledochal cyst without a remnant duct, and in anastomosis. Liuming H (2011) reported 1 case of postoperative bile leakage in a study of 39 patients [19]. In a study of Acker SN (2013), the rate of bile leakage was 1.6%. Pancreatitis was a relatively rare postoperative complication after choledochal cyst excision. Preoperative pancreatitis might be an important risk factor predisposing the patient to postoperative pancreatitis [20]. Compared to other studies, the rates of early complications of our study was low.

Several tips were suggested by some authors to reduce the rate of bile leakage. Firstly, electrical dissection should not be overused and over-dissecting the anterior wall of common hepatic duct should also be avoided. Secondly, good laparoscopic suture technique should be practiced. Thirdly, bowel loop with good arterial arcade with sufficient length should be chosen to construct a tension free anastomosis. Finally, the anastomosis should be carefully checked before the end of the operation. Other reported complication including incisional fluid collection, pancreatic fistulas and early adhesive intestinal obstruction was not noted in our study. We had 2 cases with anastomotic stenosis at 2 and 3 year. The

patients were rehospitalized due to intrahepatic lithiasis. Open operation with anatomotic redo resolved the problem. We could not analyzed the long-term results because number of patients who lost the follow-up.

Postoperative follow-up results after hospital discharge to 3 months were classified as good in 90.5% of children. Nine patients were lost to follow-up. In Vietnam, patients usually do not come back to the hospital unless they feel something wrong. Some patients lived too far and in rural region to contact by letters.

Our study contains some limitations. Firstly, the sample size was small and the follow-up period was short (within 3 months). Secondly, there was no control group to compare such as open surgery to see the efficiency of laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy in children.

Conclusion

Laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy was feasible and safe in children. It could significantly improve in terms of operative time, overall postoperative complication rate, and the length of hospital stay. The short-term outcomes were good in most cases.

Acknowledgments

The authors are grateful to Prof. Pham Nhu Hiep - director of Hue Central Hospital; physicians, administrative staff at Department of Pediatric and Abdominal Emergency Surgery for allowing us to undertake this research.

Conflicts of Interest

The authors declare no conflict of interest.

Author Contribution

All authors participated in the study design, data collection, and literature search. Data was analyzed by NTX. NTX and NHS wrote the paper. All authors read and approved the final manuscript.

References

- Soreide K, Korner H, Havnen J, Søreide JA (2004) Bile duct cysts in adults. *Br J Surg* 91: 1538-1548.
- Tang W, Dong K, Liu G, Cui X, Zheng S (2015) The clinical characters of congenital choledochal cysts in perinatal patients: A retrospective analysis in a single institution. *Am J Perinatol* 32: 853-858.
- Kayaalp C, Soyer V, Ersan V, Aydın C, Karagül S (2016) Laparoscopic choledochal cyst excision and Roux-en-Y hepaticojejunostomy. *Ulus Cerrahi Derg* 32: 152-154.
- Diao M, Li L, Cheng W (2011) Laparoscopic versus open Roux-en-Y hepatojejunostomy for children with choledochal cysts: Intermediate-term follow-up results. *Surg Endosc* 25: 1567-1573.
- Liu Y, Yao X, Li S, Liu W, Liu L, et al. (2014) Comparison of therapeutic effects of laparoscopic and open operation for congenital choledochal cysts in adults. *Gastroenterol Res Pract*.
- Liem NT (2013) Laparoscopic surgery for choledochal cysts. *J Hepatobiliary Pancreat Sci* 20: 487-491.
- Duan X, Mao X, Jiang B, Wu J (2015) Totally laparoscopic cyst excision and Roux-en-Y hepaticojejunostomy for choledochal cyst in adults: A single-institute experience of 5 years. *Surg Laparosc Endosc Percutan Tech* 25: e65-e68.
- Xuan NT, Hiep PN, Thien HH, Vu PA, Thanh PH, et al. (2013) Laparoscopic excision of choledochal cyst in Hue Central Hospital. *Vietnam Laparoscopic surgery and Endoscopy Journal* 1: 22-25.
- Terblanche J, Worthley CS, Spence RA, Krige JE (1990) High or low hepaticojejunostomy for bile duct strictures? *Surgery* 108: 828-834.
- Wiseman K, Buczkowski AK, Chung SW, Francoeur J, Schaeffer D, et al. (2005) Epidemiology, presentation, diagnosis, and outcomes of choledochal cysts in adults in an urban environment. *Am J Surg* 189: 527-531.
- Lee SE, Jang JY, Lee YJ, Choi DW, Lee WJ, et al. (2011) Choledochal cyst and associated malignant tumors in adults: A multicenter survey in South Korea. *Arch Surg* 146: 1178-1184.
- Choi SB, Choi SY (2016) Current status and future perspective of laparoscopic surgery in hepatobiliary disease. *Kaohsiung J Med Sci* 32: 281-291.
- Farello GA, Cerofolini A, Rebonato M, Bergamaschi G, Ferrari C, et al. (1995) Congenital choledochal cyst: Video-guided laparoscopic treatment. *Surg Laparosc Endosc* 5: 354-358.
- Sheng Q, Lv Z, Xu W, Xiao X, Liu J, et al. (2017) Reoperation after cyst excision with hepaticojejunostomy for choledochal cysts: Our experience in 18 cases. *Med Sci Monit* 23: 1371-1377.
- Huang CS, Huang CC, Chen DF (2010) Choledochal cysts: Differences between pediatric and adult patients. *J Gastrointest Surg* 14: 1105-1110.
- Liem NT, Pham HD, Dung le A, Son TN, Vu HM (2012) Early and intermediate outcomes of laparoscopic surgery for choledochal cysts with 400 patients. *J Laparoendosc Adv Surg Tech A* 22: 599-603.
- Tang ST, Yang Y, Wang Y, Mao YZ, Li SW, et al. (2011) Laparoscopic choledochal cyst excision, hepaticojejunostomy, and extracorporeal Roux-en-Y anastomosis: A technical skill and intermediate-term report in 62 cases. *Surg Endosc* 25: 416-422.
- Ono S, Fumino S, Shimadera S, Iwai N (2010) Long-term outcomes after hepaticojejunostomy for choledochal cyst: A 10- to 27-year follow-up. *J Pediatr Surg* 45: 376-378.
- Liuming H, Hongwu Z, Gang L, Jun J, Wenying H, et al. (2011) The effect of laparoscopic excision vs open excision in children with choledochal cyst: A midterm follow-up study. *J Pediatr Surg* 46: 662-665.
- Acker SN, Bruny JL, Narkewicz MR, Roach JP, Rogers A, et al. (2013) Preoperative imaging does not predict intrahepatic involvement in choledochal cysts. *J Pediatr Surg* 48: 2378-2382.