



REVIEW ARTICLE

The Role of Microbiome in Malaria Transmission and Severity

Kwame Kumi Asare^{1,2*} and Paul Ekow Duntu²

¹Department of Biomedical Sciences, School of Allied Health Sciences, College of Allied Health Sciences, University of Cape Coast, Cape Coast, Ghana

²Biomedical and Clinical Research Centre, College of Allied Health Sciences, University of Cape Coast, Cape Coast, Ghana

*Corresponding author: Kwame Kumi Asare, Department of Biomedical Sciences, Biomedical and Clinical Research Centre, School of Allied Health Sciences, College of Allied Health Sciences, University of Cape Coast, Cape Coast, Ghana



Abstract

Malaria is a public health concern, especially in sub-Saharan Africa. Several malaria control programs to treat, control, and eradicate malaria have been implemented. However, more than 90% of malaria-related deaths in under five years occur in Africa. With the emergence and spread of artemisinin-resistant *P. falciparum*, global malaria control now focuses on vaccine development and a search for novel and efficient antimalarial drugs. Although current evidence shows host microbiota play a significant role in malaria transmission and disease severity, little attention has focused on advancing the understanding of the role played by microbiome in malaria infection and the pathogenesis of the disease. This review discusses the interplay between malaria and host microbiomes and their role in malaria transmission and disease severity. Microbiota is known to regulate malaria survival and fertility, control transmission and modulate immunity against malaria infection. Thus, advancing the study of the interaction between malaria infection and microbiome could broaden the therapeutic advancement and the discovery of new and potent antimalarial drugs to tackle the various aspects of malaria infection and block malaria transmissions.

Keywords

Plasmodium falciparum, Microbiota, Malaria transmission, Disease severity, Novel antimalarial drug discovery, Malaria immunity, Immune escape

Introduction

Malaria is a parasitic disease that affects humans globally [1]. Five causative organisms, *P. falciparum*, *P. malariae*, *P. ovale*, *P. vivax* and *P. knowlesi*, infect

humans. Malaria kills over 660,000 people annually, mostly children under five years [2]. *P. falciparum* is the most virulent malaria parasite and is predominantly transmitted by the *Anopheles gambiae* vector in sub-Saharan Africa [1,3-5].

The transmission of parasites across vertebrate and invertebrate hosts involves a series of complex interactions that allow the parasite to survive, invade, and replicate in different cells and tissues. Mosquitoes ingest sexual forms of malaria parasites (macrogametocytes and microgametocytes) that fuse to form gametes in the midgut. The gametes develop into sporozoites that invade the salivary glands. Malaria parasites adapt to the host environmental niche inhabited by a community of microbes (microbiota). The host microbiota plays a crucial role in the pathophysiology of invading pathogens (Table 1). The symbiotic interactions between microbiota and pathogenic organisms contribute to the acquisition of disease resistance or abiotic stress tolerance in both invertebrate and vertebrate hosts [6]. The microbiota composition of the mosquito's midgut and salivary glands determines the capacity of a mosquito to transmit pathogens and is affected by the mosquito's immunity, longevity, fertility, and metabolism [7-10]. Bacteria inhabit almost every part of the female *Anopheles* mosquito, the vector of malaria [11-13]. Bacteria-malaria parasite interactions either promote or inhibit survival and growth in the midguts of vectors [14,15].



Citation: Asare KK, Duntu PE (2024) The Role of Microbiome in Malaria Transmission and Severity. J Infect Dis Epidemiol 10:324. doi.org/10.23937/2474-3658/1510324

Accepted: July 08, 2024; Published: July 10, 2024

Copyright: © 2024 Asare KK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 1: Common microbes involved in vector and host microbiota and infected *Plasmodium* species.

	Vector microbiota		Host microbiota	
	Inhibition	No Inhibition	Skin	Gut
<i>P. falciparum</i>	<p>General microbiota: <i>Pseudomonadaceae</i>, <i>Enterobacteriaceae</i>, <i>Aeromonadaceae</i>, <i>Comamonadaceae</i>, <i>Moraxellaceae</i>, <i>Cyanobacteria</i>, <i>Serratia</i></p> <p>Vector specific microbiota:</p> <p><i>An. Stephensi</i>: <i>Escherichia coli</i> H243; <i>Pseudomonas aeruginosa</i>; <i>Ewingella americana</i>; <i>E. coli</i> HS5; <i>P. aeruginosa</i>; <i>Serratia marcescens</i>; <i>Xanthomonas malthophila</i>; <i>Cedecea lapagei</i></p> <p><i>An. gambiae</i>: <i>S. aureus</i>, <i>E. coli</i>, <i>Enterobacter</i> sp., <i>Chromobacterium</i> sp., <i>P. putida</i>; <i>Pantoea</i> sp.; <i>S. marcescens</i>, <i>Serratia</i>; <i>Methylobacterium</i></p> <p><i>An. coluzzii</i>: <i>E. coli</i>; <i>S. marcescens</i>; <i>Pseudomonas sutzeri</i>; <i>Comamonas</i> spp.; <i>Enterobacter</i> spp.; <i>B. pumilus</i></p>	<p><i>An. Stephensi</i>: <i>Staphylococcus aureus</i>; <i>S. epidermidis</i>; <i>E. coli</i> HB101</p> <p><i>An. gambiae</i>: <i>Bacillus pumilus</i></p> <p><i>An. coluzzii</i>: <i>Acinetobacter septicus</i></p>	<p><i>Corynebacteria</i>, <i>Staphylococcus</i>, <i>Propionibacterium</i>, <i>Micrococcus</i>, <i>Pseudomonas</i>, <i>Brevibacterium</i> species, <i>Pityrosporum</i>, <i>Pityrosporum</i></p>	<p><i>Enterobacter</i>, <i>Enterococci</i>, <i>Bifidobacterium</i>, <i>Bacteroides</i>, <i>Clostridium</i>, <i>Firmicute</i>, <i>Bacteroidetes</i>, <i>Proteobacteria</i>, <i>Actinobacteria</i>, <i>Verrucomicrobia</i></p>
<i>P. vivax</i>	<p><i>Enterobacter</i>, <i>Serratia</i></p> <p><i>An. albimanus</i>: <i>S. marcescens</i>; <i>Enterobacter cloacae</i>; <i>Enterobacter amnigenus</i></p>			
<i>P. berghei</i>	<p><i>An. gambiae</i>, and <i>An. coluzzii</i>: <i>S. aureus</i>; <i>E. coli</i>; <i>E. cloacae</i></p> <p><i>An. stephensi</i>: <i>S. marcescens</i> HB3</p>	<p><i>An. stephensi</i>: <i>S. marcescens</i> HB18</p>		
<i>P. yoelii</i>	<p><i>An. gambiae</i>: <i>Pseudomonas aeruginosa</i>, <i>Ps. vesicularis</i> and <i>Cedecea lapagei</i>.</p>			

The mosquito feeding patterns, risks, and severity of malaria infection are affected by the human microbiota [16-20]. The recognition that malaria parasites interact with the host microbes to establish infection and virulence at various stages of the parasite life cycle has attracted the interest of researchers and clinicians. Understanding the specific role played by the host microbiome in malaria transmission and pathogenesis is critical for developing strategies for the prevention, control, and eradication of malaria globally. The review discusses the significance of microbiota in malaria transmission and severity.

The Mosquito Gut Microbiota and Malaria

Microbiota may have several metabolic effects that are often detrimental to invading pathogens. However, changes in the microbiota composition may increase susceptibility to infectious and non-infectious diseases [21]. The clinical manifestations of pathogenic infections

may be affected by the host microbiota and the host immune system [22]. Microbiota modulates pathogenic infections by augmenting the innate and adaptive immune systems [23] and plays a variety of functions ranging from food supplementation, improvement of digestive systems, toleration of environmental disturbances, protection against parasites, and enhancement and maintenance of homeostasis of the host immune system [24]. In the mosquito vector, the microbiotas play a role in nutrition, development, and reproduction [4], such that a change or removal in microbiota composition in mosquitoes could reduce fitness and the phenotype of the mosquito host [25].

Bacteria are the most prominent microbiota found in mosquitoes and are predominantly found in the digestive tract of mosquitoes. *Acetobacteriaceae* are the most dominant bacteria that affect both wild and laboratory-raised colonies of Anopheline and Aedes mosquitoes [26].

The mosquito midgut harbours digestive enzymes and an innate immune system that inhibit disease transmission [27,28], by influencing the susceptibility of the insect to *Plasmodium* infection [29-32]. Although the microbiota might protect mosquitoes from pathogenic infections [10,33], organisms such as viruses, helminths, and protozoans have devised ways to take advantage of the gut microbiota to spread [34,35]. The *Enterobacteriaceae* and *P. falciparum* infections in mosquitoes suggest possible bacteria-parasite interactions playing a role in malaria parasite development [10,33]. Both laboratory-cultivated and field-collected mosquitoes harbour varieties of bacteria such as *Asaia*, *Enterobacter*, *Mycobacterium*, *Sphingomonas*, *Serratia*, and *Chryseobacter* in their midgut [36,37].

Variations in Mosquito Gut Microbiota

Mosquitoes harbour a diverse microbial flora that contributes to the expansion of the mosquito gene pool [33]. The mosquito-microbiota genotype interaction influences the variations in specific immunological phenotypes and the adaptation of gut microbes. The mosquito gut microbiota comprises prokaryotes, viruses, and eukaryotic microorganisms that shape diversity in vector competence across diverse mosquito-borne diseases and pathogenic species. *Bacteroidetes*, *Proteobacteria*, *Actinobacteria*, and *Firmicutes* are the major microbiota in the midgut of adult mosquitoes [38]. Mosquito populations in different geographical settings exhibit different levels of parasitic diseases or parasite-specific infections, suggesting a possible role of inhabited microbiota in the transmission of parasites [34,39]. The presence of interspecific rivalry, the generation of poisons and inhibitory substances. And specific bacterial taxa may render the mosquito midgut inhospitable for some bacteria or malaria parasites. Mosquitoes bred in different locations exhibit diverse midgut bacteria compositions suggestive of the environmental role in gut microbiota, nutritional status, developmental stages, and mosquito species [10]. Mosquito gut microbiota may have been influenced by the consumption of aquatic bacteria by larvae, which is alkaline (pH range of 8 to 11 & high quantities of carbonate ions) and favours the establishment of a subset of bacterial infections, similar to humans [40-42]. Factors such as redox potential, specific nutrients and proteolytic enzymes in the gut, early colonists, and selective pressures determine the mosquito gut microbiota [43]. The novel bacterial community shaped by maternally transmitted germs may compete with the new colonists. For instance, maternally inherited bacteria compete with *Asaia* in the reproductive tract of the Asian malaria vector, *Anopheles stephensi* [44-48]. Although all mosquito species typically consume blood and nectar, some mosquito species exhibit marked variations in their preferred larval habitats and

sugar, and blood meal hosts, which may pre-expose them to different microbes. This may partially explain the observed differences. The midgut microbiota may enhance parasite infection, showing that *Plasmodium falciparum* development in the *Anopheles gambiae* mosquito depends on complex vector-microbes-parasite interactions [33].

Effects of Mosquito Gut Microbiota on Malaria Transmission

Transmission of malaria depends on the success of the several transitional stages that the malaria parasites go through in the midgut during a series of intricate developmental stages [49,50]. The development from motile ookinetes to oocysts migrates through the midgut epithelium; most oocysts die when resting under the basal lamina of the midgut wall [50]. The microbiota could influence the parasite's development and viability, from the ookinete to the formation of oocysts and penetrating midgut epithelial cells [24]. The gut microbiota and the metabolic by-products determine the pathogenic effects of a compromised gut barrier [51]. The microbiota decreases mosquito vulnerability to *Plasmodium* infections by stimulating mosquito immunity, forming a physical barrier, and producing reactive oxygen species to inhibit *Plasmodium* infection [9,15]. There are variations in the vulnerability of wild *Anopheles* populations, and these observed variations are associated with the presence or absence of specific gut microbiota [9]. Also, only a few fractions of the natural *Anopheles* population carry specific gut microbiota [9]. The *Anopheles gambiae* and *Anopheles funestus* in the field harbour 16 bacterial species among the 14 genera [52]. The ability of microbiota to control the malaria parasite infections as well as alter the mosquito lifespan suggests that the gut microbiome could prevent malaria transmission [37,53,54]. Gram-negative bacteria such as *Enterobacter Esp Z*, *Chromobacterium Csp*, and *Serratia marcescens* in the midgut of *Anopheles* mosquitoes inhibit *P. falciparum* development [9,55-57]. Furthermore, *Wickerhamomyces anomalous*, a yeast from the midgut of *Anopheles stephensi*, produces a lethal toxin that suppresses *P. berghei* ookinetes in the presence of 1-3-glucanase *in vitro* [58]. Also, chitinase secreted by *Plasmodium* ookinetes allows the crossing of the peritrophic matrix [57,59]. Thus, barriers or bacterial toxins may put selection pressure on malaria parasites.

The Human Microbiota

Humans harbour complex communities of symbiotic microbes such as bacteria, archaea, viruses, and fungi. The gastrointestinal microbiome contains roughly millions of distinct genes, mostly from bacteria. Factors such as gestational age, diet, antibiotic exposure, probiotic usage and nutritional supplements, hygienic conditions, host genetics, and host immune system

could influence the microbiota. The human microbiome is essential for the immune system, the brain, nutrition, and metabolism [60]. Antimicrobial drugs such as antimalarials or antibiotics may poison gut microbial ecosystems and the normal physiological function of the microbiota [61,62].

The Human Gut Microbiota and *Plasmodium* Infections

The gastrointestinal microbiota constitutes about 70% of all human microbiota. The bacterial population is diverse, has been associated with several disease conditions, such as malnourishment and allergies, and has a severe impact on cognitive and physical development [63-65]. *Enterobacter*, *Enterococci*, *Bifidobacterium*, *Bacteroides*, *Clostridium*, *Firmicute*, *Bacteroidetes*, *Proteobacteria*, *Actinobacteria*, and *Verrucomicrobia* dominate the gut microbiota [59,66-68]. The small intestines contain monosaccharides, disaccharides, and amino acids and harbour *proteobacteria*, *Lactobacillales*, *Enterococcus*, and *Akkermansia*, and the lumen harbours *Enterobacteriaceae* and *Bacteroides*, *Bifidobacterium*, *Streptococcus*, *Enterococcus*, *Clostridium*, *Lactobacillus*, and *Ruminococcus* [67,69]. The gut microbiota also includes viruses such as bacteriophages (Picornaviridae, Reoviridae, Astroviridae and Caliciviridae) which have a nonpathogenic relationship with the human host [70]. The bacteriophages produce lysogenic and lytic substances for prevalent viral infections in healthy human tissues [70].

The gut microbiota, environmental factors, and genetic variations are associated with variabilities in the clinical episodes of *Plasmodium* infections [71]. Frequently, *Bifidobacterium* and *Streptococcus* species isolated from the guts of individuals with no *P. falciparum* infections, are known to offer resistance to malaria infection. Similarly, *Lactobacillus* and *Bifidobacteria* species protect mice from *P. yoelii* infection. Their protective effects are, however, affected by diet alterations and probiotic administration [23]. The gut microbiota differentially expresses toxic metabolites in severe malaria in human and animal models that could be detrimental to malaria parasites [72,73]. Blood-stage malaria parasites are more vulnerable to metabolic dysregulation driven by either antimalarials or the microbiota [74].

Malaria and Mammalian Gut Microbiota

Microbiota and bacterial communities influence the severity of malaria [75]. The cardinal symptoms of diarrhoea and abdominal discomforts associated with *Plasmodium falciparum* infection result from pathological alteration of the gut [76]. Malaria infections cause changes in villi and lead to bleeding of mucosal tissues due to rupturing [77]. Human cases of severe *falciparum* malaria increase intestinal permeability [78]. This alteration of the gut environment influences the

progression of malaria infection [76]. The severity and disease progression of *Plasmodium* infection in C57BL/6 and BALB/c mice coincide with the extent of alteration in gut microbiota [76]. *P. yoelii* 17XNL and *Heligmosomoides polygyrus* coinfection protect mice from fatal malaria infections [79]. Tomoyo and associates have shown that severe intestinal pathology during malaria is associated with changes in the microbiota [76]. Different bacterial communities modulate malaria infections and disease severity differently. For instance, *Lactobacillus* reduces parasitemia and causes milder cerebral malaria [76]. Hence, the vulnerability of individuals to *Plasmodium* infections, disease subtypes, and mosquito affinity is related to gut microbiome dysbiosis [16,80,81]. Although variations in immune response and host genetics play a role in parasite defense, the current information shows that gut microbiota significantly affects the transmission and development of malaria [22,82].

The Skin Microbiota and Mosquito Attraction

The skin forms the first line of defense against environmental and pathogenic exposures [83]. The skin microbiota varies among individuals and is affected by individual behaviour, genetic makeup, age, sex, and environmental factors [84-87]. These factors indirectly influence the choice of mosquitoes for their prospective human hosts for disease transmission [85]. Individuals produce volatile odour substances [88] which are modified by the skin microbiota and vary among individuals and their exposure to mosquito vectors. The skin microbiota produces odour by breaking down substances into by-products such as methyl ketones, carboxylic acids, and linear alcohols, and these odours serve as chemical signals to draw mosquito vectors to the host [83,89,90]. The strength of odour is associated with specific bacterial compositions [90-96]. Female mosquitoes use unique olfactory profiles to detect host-derived chemical and physical signals (such as heat and visual cues) to determine hosts of choice [16,97-99]. The vector-host feeding interaction influences the transmission of the disease [100,101].

The various skin microbes have unique metabolic pathways and by-products [83], and early studies showed that *Corynebacteria* distinctively produced apocrine odour from volatile fatty acids, while *Staphylococcus* species convert branched-chain amino acid residues into *short-chain residues*, which are extremely odorous and volatile [102]. Anthropophilic anopheline mosquitoes that transmit malaria to humans exhibit variations in their attraction to different individuals [88]. The unique composition of the skin microbiota impacts the distinctive odour profiles of individuals [88]. A previous study reported that female *Anopheles gambiae sensu stricto* is attracted to sweat samples incubated under aerobic conditions [80]. Similarly, female *Anopheles gambiae* is more attracted to lures made from skin microorganisms from the feet of

healthy individuals than the control traps made of sterile agar and clean air with or without carbon dioxide [81]. *Corynebacterium*, *Staphylococcus*, *Propionibacterium*, *Micrococcus*, *Pseudomonas*, *Brevibacterium* species and *Pityrosporum* species are possible skin microbes that produce chemoattractants. *Pseudomonas aeruginosa* and *Staphylococcus epidermis* are essential bacteria species that determine the attractiveness of mosquitoes to the host [80,103]. Verhulst et al. showed that people who are attracted to mosquitoes have a less varied microbiome [89]. Also, individuals with *Pseudomonas* poorly attract mosquitoes [83]. This pattern of intraspecies variations in mosquito attraction is essential for pathogen exposure and transmission. Thus, individuals that attract mosquitoes more frequently have a high risk of contracting a pathogen carried by a mosquito and promoting its spreading [83].

Microbiota and Immunity in Malaria

Microbiota interacts with malaria infections by producing inhibitory bioactive toxins and enzymes or stimulating the host immune system [24]. Surface glycan Gal-13 Gal-4 GlcNAc (-galactosyl) expressed by *Enterobacteriaceae* such as *Klebsiella*, *Serratia*, and *Escherichia* induces the host to produce anti-galactosyl antibodies that interact with *Plasmodium* sporozoites and inhibit hepatic invasion [104,105]. A study revealed that Malians with high anti-galactosyl immunoglobulin M (IgM) have a low occurrence of *P. falciparum* infection [104]. *Streptococcus*, *Bifidobacterium*, *Escherichia* and *Shigella* provide cross-protection against *P. falciparum* infection [82]. Microbe-stimulated antibodies opsonize *Plasmodium*-parasitized erythrocytes and process them for spleen clearance [75]. Also, the recruitment of phagocytic cells by *Bifidobacteriaceae* could regulate outcomes in malaria [106,107]. The peptidoglycan recognition protein (PGRP) family functions as pattern recognition receptors and effectors for innate immunity and regulates microbial growth and parasite infection in mosquitoes [108]. The host immune response to blood-stage malaria parasites involves RBC sequestration and systemic-induced IL-1 β , IL-6, IL-8, IL-12p70, interferon (IFN)- γ , and tumour necrosis factor- α (TNF)- α pro-inflammatory cytokines and chemokines (e.g., interleukin IL-1 β , IL-6, IL-8, IL-12p70, interferon (IFN)- γ , and tumour necrosis factor- α (TNF)- α) and innate immune cells such as natural killers (NKs), mast cells, neutrophils, and antibodies. Previous studies showed that the lack of interleukin-21 (IL-21) signalling within the spleen leads to impaired clearance of *Plasmodium* infection. T follicular helper (Tfh) and germinal centre (GC) B cells are critical in developing immune antibodies against *Plasmodium* species.

The disruption of IL-21 inhibits the production of immunity by Tfh cells, leading to sustained high parasitemia. The Tfh cell disruption in mice infected with *P. yoelii* is able to self-resolve chronic *P. chabaudi*

infection. Impaired Tfh differentiation and inefficient GC responses cause severe malaria infection. IFN γ and IL-10 cytokines affect the humoral immune response to *Plasmodium* and control parasite control and host survival. The spleen germinal centre (GC) activation and coordination of humoral immunity are responsible for clearing *Plasmodium* infection. The clinical presentation of malaria depends on a balance between pro-inflammatory and regulatory cytokines such as IL-10, transforming growth factor beta (TGF β), and regulatory T lymphocytes, IL-10-producing T lymphocytes immune cells.

The microbiota prevents malaria transmission by inducing anti- α -gal IgM and IgG (IgG2b, IgG3) in the skin to employ an immune complement pathway to inhibit liver invasion and sporozoites transmission. CD8+, CD4+, and T cells in the skin and the liver are adaptive immune factors that target antigens specific to intracellular pathogens and act as initial sterilizing immunity. IFN- γ , TNF, and IL-2 inhibit the early stages of malaria parasites and protect against sporozoites and blood-stage antigens. The NK, NKT, $\gamma\delta$ T cells and macrophages induced by microbiota in the peripheral blood circulation stimulate nitric oxide and nitric oxide synthase. The gut microbiota downregulates IL-1 β , IL-6, IL-8, and IL-12 to control blood-stage malaria parasites. TGF- β and IL-10 regulatory cytokines reduce malaria parasitaemia and Treg, DCs balance the immune system and present parasites to the T cells and the activation of B cells for parasite clearance.

The cross-talk of host-microbiota-malaria immunity can modulate the risk of *Plasmodium* infection, transmission and disease severity. Understanding the immune interaction between the host-microbiota and malaria infection could give insight into an innovative approach to preventing and managing malaria in endemic countries.

Malaria, Antibiotics, and the Microbiome

Sub-optimal antibiotics treatment of microbiota promotes dysbiosis and affects mosquito gut bacteria [109-111]. Notably, the addition of a cocktail of streptomycin and penicillin to mosquito blood meals enhanced insect survival, fertility, and toleration of *Plasmodium* infection [112]. Exposure of high doses of doxycycline to gut microflora resulted in increased *P. falciparum* infection load, similar to the effect of penicillin-streptomycin [112,113]. However, azithromycin or co-trimoxazole treatment is detrimental to both midgut microbiota and the *Plasmodium* infection. Thus, characterizing antibiotic usage could be a novel method for controlling malaria transmission.

A previous study suggested that malaria may predispose to bacteraemia as acute *P. falciparum* malaria causes widespread immunosuppression and decreases macrophage functionality to phagocytosis-

infected parasites [114,115].

Future Perspective

Mosquito-transmitted epidemics like malaria, yellow fever, dengue fever, chikungunya fever and Zika fever cause roughly 350 million illnesses and 500000 fatalities worldwide each year [116]. Considering that the majority of these diseases lack an effective vaccination [117], one of the most effective methods to prevent disease is vector control [118]. Rapid scientific advancement in the study of the microbiome has invented different therapeutic possibilities and revived "primitive techniques" that target the microbiota and may be useful in the fight against infectious diseases. The impact of microbiota and invasive pathogens on mosquito capacity to transmit malaria parasites in contrast to the parasite itself provide new approaches for malaria biocontrol [119]. The application of microbes to prevent the spread of malaria has drawn attention worldwide. A high-tech method based on genetically altered mosquitoes has several advantages over a low-tech one when used as a control strategy. These include mass manufacturing of *in vitro* cultures, the ability to store and transfer products to remote locations, the absence of evolutionary pressure, mosquito population coverage, the absence of side effects, and needless genetic change [120]. The positive outcomes of the microbiota and mosquito-derived microbiota could be used as biological agents against the *Plasmodium* species and/or its mosquito vectors. Also using synthetic microbes to modify the genetic composition of *Anopheles* vectors may help reduce the spread of malaria [119]. Entomopathogenic fungi such as *Beauveria bassiana* (Bals. -Criv.) Vuill. and *Metarhizium anisopliae* Metschn. have been proposed as biological control agents for malaria [121]. Recent research has shown the value and possibility of using permethrin and entomopathogenic fungi together as pesticide resistance prevention measures for vectors *Angambiae s.l.* [122,123].

Conclusion

Understanding the role microbiota play in the transmission, control of malaria infections (symptomatic and asymptomatic) through the modulation of vector and host immunities has become essential with the emergence and spread of artemisinin-resistant *P. falciparum*. The microbiota has been shown to affect malaria survival and fertility, as well as control transmission and alter immunity to malaria infection. Thus, furthering the study of the interaction between malaria infection and microbiome could extend therapeutic progress and the identification of novel and strong antimalarial medications to address the various aspects of malaria infection and block malaria transmissions.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All the data are available in the manuscript.

Competing interests

The authors declare no competing interests.

Funding

AKK was partly supported with the 2022/2023 Ghana government books and research allowance BRA2022/2023.

Authors' contributions

Conceptualization, KKA; Methodology, KKA, PED; Supervision, KKA; Writing – original draft, PED; Writing – review & editing; KKA, PED.

Acknowledgments

We thank Dr Yeboah Kwaku Opoku, Department of Biology Education, Faculty of Science Education, University of Education, Winneba, Ghana for reviewing the manuscript.

References

1. Health Organization W. WORLD MALARIA REPORT 2016 ISBN 978 92 4 151171 1.
2. (2021) World malaria report 2021.
3. Cumberland S (2009) Mosquito wars. Bull World Heal Organ 87: 167-168.
4. Minard G, Mavingui P, Moro CV (2013) Diversity and function of bacterial microbiota in the mosquito holobiont. Parasites and Vectors 6: 1-12.
5. Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, et al. (2018) Vital Signs: Trends in reported vectorborne disease cases - United States and Territories, 2004-2016. Morbidity Mortality Weekly Report 67: 496-501.
6. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, et al. (2009) A Wolbachia Symbiont in *Aedes aegypti* limits infection with Dengue, Chikungunya, and Plasmodium. Cell 139: 1268-1278.
7. Agustin DP, Erlambang BDP, Cahyanti CT, Waton S, Oktarianti R, et al. (2021) *Anopheles* microbiota in malaria vector and the effect on parasite transmission. Bioedukasi J Biol dan Pembelajarannya XIX: 79-89.
8. Heu K, Gendrin M (2018) Mosquito microbiota and its influence on disease vectorial transmission. Biol Aujourdhui 212: 119-136.
9. Cirimotich CM, Dong Y, Clayton AM, Sandiford SL, Souza-Neto JA, et al. (2011) Natural microbe-mediated refractoriness to Plasmodium infection in *Anopheles*

- gambiae*. *Science* 332: 855-858.
10. Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, et al. (2012) Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. *PLoS Pathog* 8: e1002742.
 11. Villegas LM, Pimenta PFP (2014) Metagenomics, paratransgenesis and the *Anopheles* microbiome: A portrait of the geographical distribution of the anopheline microbiota based on a meta-analysis of reported taxa. *Mem Inst Oswaldo Cruz* 109: 672-684.
 12. Sharma P, Sharma S, Maurya RK, Das De T, Thomas T, et al. (2014) Salivary glands harbor more diverse microbial communities than gut in *Anopheles culicifacies*. *Parasit Vectors* 7: 235.
 13. Hillesland H, Read A, Subhadra B, Hurwitz I, McKelvey R, et al. (2008) Identification of aerobic gut bacteria from the kala azar vector, *Phlebotomus argentipes*: A platform for potential paratransgenic manipulation of sand flies. *Am J Trop Med Hyg* 79: 881-886.
 14. Wang M, An Y, Gao L, Dong S, Zhou X, et al. (2021) Glucose-mediated proliferation of a gut commensal bacterium promotes *Plasmodium* infection by increasing mosquito midgut pH. *Cell Reports* 35: 108992.
 15. Bando H, Okado K, Guelbeogo WM, Badolo A, Aonuma H, et al. (2013) Intra-specific diversity of *Serratia marcescens* in *Anopheles* mosquito midgut defines *Plasmodium* transmission capacity. *Sci Reports* 3: 1641.
 16. Verhulst NO, Qiu YT, Beijleveld H, Maliepaard C, Knights D, et al. (2011) Composition of human skin microbiota affects attractiveness to malaria mosquitoes. *PLoS One* 6: e28991.
 17. Takken W, Verhulst NO (2017) Chemical signaling in mosquito-host interactions: The role of human skin microbiota. *Curr Opin Insect Sci* 20: 68-74.
 18. Mukherjee D, Chora ÂF, Mota MM (2020) Microbiota, a Third Player in the Host-*Plasmodium* Affair. *Trends Parasitol* 36: 11-18.
 19. Smith CDM, Gong M, Andrew AK, Russ BN, Ge Y, et al. (2019) Composition of the gut microbiota transcends genetic determinants of malaria infection severity and influences pregnancy outcome. *EBioMedicine* 44: 639-655.
 20. Villarino NF, LeClerc GR, Denny JE, Dearth SP, Harding CL, et al. (2016) Composition of the gut microbiota modulates the severity of malaria. *Proc Natl Acad Sci USA* 113: 2235-2240.
 21. Sirisinha S (2016) The potential impact of gut microbiota on your health: Current status and future challenges. *Asian Pac J Allergy Immunol* 34: 249-264.
 22. Bär AK, Phukan N, Pinheiro J, Simoes-Barbosa A (2015) The Interplay of host microbiota and parasitic protozoans at mucosal interfaces: Implications for the outcomes of infections and diseases. *PLoS Negl Trop Dis* 9: e0004176.
 23. Netea MG, Joosten LAB, Latz E, Mills KHG, Natoli G, et al. (2016) Trained immunity: A program of innate immune memory in health and disease. *Science* 352: aaf1098.
 24. Weiss B, Aksoy S (2011) Microbiome influences on insect host vector competence. *Trends Parasitol* 27: 514-522.
 25. Koch H, Schmid-hempel P (2011) Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *PNAS* 108: 19288-19292.
 26. Damiani C, Ricci I, Crotti E, Rossi P, Rizzi A, et al. (2010) Mosquito-bacteria symbiosis: The Case of *Anopheles gambiae* and *Asaia*. *Microb Ecol* 60: 644-654.
 27. Bennett KE, Olson KE, Muñoz M de L, Fernandez-Salas I, Farfan-Ale JA, et al. (2002) Variation in vector competence for dengue 2 virus among 24 collections of *Aedes aegypti* from Mexico and the United States. *Am J Trop Med Hyg* 67: 85-92.
 28. Xi Z, Ramirez JL, Dimopoulos G (2008) The *Aedes aegypti* toll pathway controls dengue virus infection. *PLoS Pathog* 4: e1000098.
 29. Ramirez JL, Souza-Neto J, Cosme RT, Rovira J, Ortiz A, et al. (2012) Reciprocal tripartite interactions between the *Aedes aegypti* midgut microbiota, innate immune system and dengue virus influences vector competence. *PLoS Negl Trop Dis* 6: e1561.
 30. Apte-Deshpande A, Paingankar M, Gokhale MD, Deobagkar DN (2012) *Serratia odorifera* a midgut inhabitant of *Aedes aegypti* mosquito enhances its susceptibility to dengue-2 virus. *PLoS One* 7: e40401.
 31. Cirimotich CM, Ramirez JL, Dimopoulos G (2011) Native microbiota shape insect vector competence for human pathogens minireview. *Cell Host Microbe* 10: 307-310.
 32. Djadid ND, Jazayeri H, Raz A, Favia G, Ricci I, et al. (2011) Identification of the Midgut Microbiota of *An. stephensi* and *An. maculipennis* for their application as a paratransgenic tool against Malaria *PLoS One* 6: e28484.
 33. Tchioffo MT, Boissière A, Churcher TS, Abate L, Gimonneau G, et al. (2013) Modulation of malaria infection in *Anopheles gambiae* mosquitoes exposed to natural midgut bacteria. *PLoS One* 8: e81663.
 34. Leibold MA, Storch D, Gaston KJ, Davies KF, Stafford HD, et al. (2010) Exploitation of the intestinal microflora by the parasitic nematode *Trichuris muris*. 328: 1391-1394.
 35. Kane M, Case LK, Kopaskie K, Kozlova A, MacDermid C, et al. (2011) Successful transmission of a retrovirus depends on the commensal microbiota. *Science* 334: 245-249.
 36. Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, et al. (2007) Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc Natl Acad Sci USA* 104: 9047-9051.
 37. Dong Y, Manfredini F, Dimopoulos G (2009) Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog* 5: e1000423.
 38. Geib SM, Jimenez-gasco MDM, Carlson JE, Tien M, Hoover K (2009) Effect of host tree species on cellulase activity and bacterial community composition in the gut of larval Asian Longhorned beetle. *Environ Entomol* 38: 686-699.
 39. Ndiath MO, Cohuet A, Gaye A, Konate L, Mazonot C, et al. (2011) Comparative susceptibility to *Plasmodium falciparum* of the molecular forms M and S of *Anopheles gambiae* and *Anopheles arabiensis*. *Malar J* 10: 269.
 40. Corena MP, Vanekeris L, Salazar MI, Bowers D, Fiedler MM, et al. (2005) Carbonic anhydrase in the adult mosquito midgut. *J Exp Biol* 208: 3263-3273.
 41. Corena MP, Fiedler MM, Vanekeris L, Tu C, Silverman DN, et al. (2004) Alkalization of larval mosquito midgut and the role of carbonic anhydrase in different species of mosquitoes. *Comp Biochem Physiol C Toxicol Pharmacol* 137: 207-225.
 42. Dadd RH (1975) Alkalinity within the midgut of mosquito larvae with alkaline-active digestive enzymes. *J Insect*

- Physiol 21: 1847-1853.
43. Dillon RJ, Dillon VM (2004) The gut bacteria of insects : Nonpathogenic interactions. *Annu Rev Entomol*.
44. Rossi P, Ricci I, Cappelli A, Damiani C, Ulissi U, et al. (2015) Mutual exclusion of *Asaia* and *Wolbachia* in the reproductive organs of mosquito vectors. *Parasit Vectors* 8: 278.
45. Jones JC (1960) The anatomy and rhythmical activities of the alimentary canal of *Anopheles* Larvae. *Annals of the Entomological Society of America* 53: 459-474.
46. Howland LJ (1930) The nutrition of mosquito larvae, with special reference to their algal food. *Bull Entomol Res* 21: 431-439.
47. Vázquez-Martínez MG, Rodríguez MH, Arredondo-Jiménez JI, Méndez-Sánchez JD, Bond-Compeán JG, et al. (2002) Cyanobacteria Associated with *Anopheles albimanus* (Diptera: Culicidae) larval habitats in Southern Mexico. *J Med Entomol* 39: 825-832.
48. Demaio J, Pumpuni CB, Kent M, Beier JC (1996) The midgut bacterial flora of wild *Aedes triseriatus*, *Culex pipiens*, and *Psorophora columbiae* mosquitoes. *Am J Trop Med Hyg* 54: 219-223.
49. Sinden RE, Dawes EJ, Alavi Y, Waldock J, Finney O, et al. (2007) Progression of *Plasmodium berghei* through *Anopheles stephensi* is density-dependent. *PLOS Pathog* 3: e195.
50. Ghosh A, Edwards MJ, Jacobs-Lorena M (2000) The Journey of the malaria parasite in the mosquito: Hopes for the New Century. *Parasitol Today* 16: 196-201.
51. Leopold S, Ghose A, Hugenholtz F, Kingston HW, Plewes K, et al. (2020) Gut microbiota composition is associated with severity of disease in patients with *P. falciparum* malaria 1-35.
52. Lindh JM, Terenius O, Faye I (2005) 16S rRNA gene-based identification of midgut bacteria from field-caught *Anopheles gambiae* Sensu Lato and *A. funestus* mosquitoes reveals new species related to known insect symbionts. *Appl Environ Microbiol* 71: 7217-7223.
53. Gonzales-Ceron L, Santillan F, Rodriguez MH, Mendez D, Hernandez-Avila JE (2003) Bacteria in midguts of field-collected *Anopheles albimanus* block *Plasmodium vivax* Sporogonic Development. *J Med Entomol* 40: 371-374.
54. Wang S, Dos-Santos ALA, Huang W, Liu KC, Oshaghi MA, et al. (2017) Driving mosquito refractoriness to *Plasmodium falciparum* with engineered symbiotic bacteria. *Science* 357: 1399-1402.
55. Dennison NJ, Saraiva RG, Cirimotich CM, Mlambo G, Mongodin EF, et al. (2016) Functional genomic analyses of *Enterobacter*, *Anopheles* and *Plasmodium* reciprocal interactions that impact vector competence. *Malar J* 15: 425.
56. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, et al. (2014) Exploring *Anopheles* gut bacteria for *Plasmodium* blocking activity. *Environ Microbiol* 16: 2980-2994.
57. Tsai YL, Hayward RE, Langer RC, Fidock DA, Vinetz JM (2001) Disruption of *Plasmodium falciparum* chitinase markedly impairs parasite invasion of mosquito midgut. *Infect Immun* 69: 4048-4054.
58. Valzano M, Cecarini V, Cappelli A, Capone A, Bozic J, et al. (2016) A yeast strain associated to *Anopheles* mosquitoes produces a toxin able to kill malaria parasites. *Malar J* 15: 21.
59. Huber M, Cabib E, Miller LH (1991) Malaria parasite chitinase and penetration of the mosquito peritrophic membrane. *Proc Natl Acad Sci USA* 88: 2807-2810.
60. Gilbert J, Blaser MJ, Caporaso JG, Jansson J, Lynch SV, et al. (2018) Current understanding of the human microbiome. *Nat Med* 24: 392-400.
61. Holmes E, Kinross J, Gibson GR, Burcelin R, Jia W, et al. (2012) Therapeutic Modulation of Microbiota-Host Metabolic Interactions. *Sci Transl Med* 4: 137rv6.
62. Appalasaamy S, Lo KY, Ch'Ng SJ, Nornadia K, Othman AS, et al. (2014) Antimicrobial activity of artemisinin and precursor derived from in vitro plantlets of *Artemisia annua* L. *Biomed Res Int* 2014: 215872.
63. Gordon JI, Dewey KG, Mills DA, Medzhitov RM (2012) The Human Gut Microbiota and Undernutrition. *Sci Transl Med* 4: 137ps12.
64. Fujimura KE, Lynch SV (2015) Microbiota in allergy and asthma and the emerging relationship with the gut microbiome. *Cell Host Microbe* 17: 592-602.
65. Mayer EA, Knight R, Mazmanian SK, Cryan JF, Tillisch K (2014) Gut Microbes and the Brain: Paradigm Shift in Neuroscience. *J Neurosci* 34: 15490-15496.
66. Hugon P, Dufour JC, Colson P, Fournier PE, Sallah K, et al. (2015) A comprehensive repertoire of prokaryotic species identified in human beings. *Lancet Infect Dis* 15: 1211-1219.
67. Bäckhed F, Fraser CM, Ringel Y, Sanders ME, Sartor RB, et al. (2012) Defining a healthy human gut microbiome: Current concepts, future directions, and clinical applications. *Cell Host Microbe* 12: 611-622.
68. Romoli O, Gendrin M (2018) The tripartite interactions between the mosquito, its microbiota and *Plasmodium*. *Parasit Vectors* 11: 200.
69. Swidsinski A, Loening-Baucke V, Lochs H, Hale LP (2005) Spatial organization of bacterial flora in normal and inflamed intestine: A fluorescence in situ hybridization study in mice. *World J Gastroenterol* 11: 1131-1140.
70. Dalmasso M, Hill C, Ross RP (2014) Exploiting gut bacteriophages for human health. *Trends Microbiol* 22: 399-405.
71. Buffie CG, Pamer EG (2013) Microbiota-mediated colonization resistance against intestinal pathogens. *Nat Rev Immunol* 13: 790-801.
72. Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyötyläinen T, Nielsen T, et al. (2016) Human gut microbes impact host serum metabolome and insulin sensitivity. *Nature* 535: 376-381.
73. Swann JR, Want EJ, Geier FM, Spagou K, Wilson ID, et al. (2011) Systemic gut microbial modulation of bile acid metabolism in host tissue compartments. *Proc Natl Acad Sci USA* 108: 4523-4530.
74. Cobbold SA, Chua HH, Nijagal B, Creek DJ, Ralph SA, et al. (2016) Metabolic Dysregulation Induced in *Plasmodium falciparum* by Dihydroartemisinin and Other Front-Line Antimalarial Drugs. *J Infect Dis* 213: 276-286.
75. Ippolito MM, Denny JE, Langelier C, Sears CL, Schmidt NW (2018) Malaria and the microbiome: A systematic review. *Clin Infect Dis* 67: 1831-1839.
76. Taniguchi T, Miyauchi E, Nakamura S, Hirai M, Suzue K, et al. (2015) *Plasmodium berghei* ANKA causes intestinal malaria associated with dysbiosis *Sci Rep* 27: 15699.

77. Olsson RA, Johnston EH (1969) Histopathologic changes and small-bowel absorption in falciparum malaria. *Am J Trop Med Hyg* 18: 355-359.
78. Wilairatana P, Meddings JB, Ho M, Vannaphan S, Looareesuwan S (1997) Increased gastrointestinal permeability in patients with Plasmodium falciparum malaria. *Clin Infect Dis* 24: 430-435.
79. Tetsutani K, Ishiwata K, Ishida H, Tu L, Torii M, et al. (2009) Concurrent infection with Heligmosomoides polygyrus suppresses anti-Plasmodium yoelii protection partially by induction of CD4 + CD25 + Foxp3 + Treg in mice. *European Journal of Immunology* 39: 2822-2830.
80. Braks MAH, Takken W (1999) Incubated human sweat but not fresh sweat attracts the malaria mosquito Anopheles gambiae sensu stricto. *Journal of Chemical Ecology* 25: 663-672.
81. Verhulst NO, Beijleveld H, Knols BG, Takken W, Schraa G, et al. (2009) Cultured skin microbiota attracts malaria mosquitoes. *Malar J* 8: 302.
82. Yooseph S, Kirkness EF, Tran TM, Harkins DM, Jones MB, et al. (2015) Stool microbiota composition is associated with the prospective risk of Plasmodium falciparum infection. *BMC Genomics* 16: 631.
83. Ruiz-López MJ (2020) Mosquito behavior and vertebrate microbiota interaction: Implications for Pathogen Transmission. *Front Microbiol* 11: 573371.
84. Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, et al. (2012) Structure, function and diversity of the healthy human microbiome. *Nat* 486: 207-214.
85. Grice EA, Kong HH, Renaud G, Young AC, NISC Comparative Sequencing Program, et al. (2008) A diversity profile of the human skin microbiota. *Genome Res* 18: 1043-1050.
86. Fierer N, Hamady M, Lauber CL, Knight R (2008) The influence of sex, handedness, and washing on the diversity of hand surface bacteria. *Proc Natl Acad Sci USA* 105: 17994-17999.
87. Leclaire S, Nielsen JF, Drea CM (2014) Bacterial communities in meerkat anal scent secretions vary with host sex, age, and group membership. *Behavioral Ecology* 25: 996-1004.
88. Leclaire S, Strandh M, Dell'Ariccia G, Gabirot M, Westerdahl H, et al. (2019) Plumage microbiota covaries with the major histocompatibility complex in blue petrels. *Mol Ecol* 28: 833-846.
89. Verhulst NO, Mukabana WR, Takken W, Smallegange RC (2011) Human skin microbiota and their volatiles as odour baits for the malaria mosquito Anopheles gambiae s.s. *Entomol Exp Appl* 139: 170-179.
90. Shelley WB, Hurley HJ, Nichols AC (1953) Axillary odor; Experimental Study of the Role of Bacteria, Apocrine Sweat, and Deodorants. *AMA Arch Derm Syphilol* 68: 430-446.
91. Leyden JJ, McGinley KJ, Holzle E, Labows JN, Kligman AM (1981) The microbiology of the human axilla and its relationship to axillary odor. *J Invest Dermatol* 77: 413-416.
92. Jackman PJ, Noble WC (1983) Normal axillary skin microflora in various populations. *Clin Exp Dermatol* 8: 259-268.
93. Kearney JN, Harnby D, Gowland G, Holland KT (1984) The follicular distribution and abundance of resident bacteria on human skin. *J Gen Microbiol* 130: 797-801.
94. Taylor D, Daulby A, Grimshaw S, James G, Mercer J, et al. (2003) Characterization of the microflora of the human axilla. *Int J Cosmet Sci* 25: 137-145.
95. Ara K, Hama M, Akiba S, Koike K, Okisaka K, et al. (2006) Foot odor due to microbial metabolism and its control. *Can J Microbiol* 52: 357-364.
96. Olang EA, Okal MN, Mbadi PA, Kokwaro ED, Mukabana WR (2010) Attraction of Anopheles gambiae to odour baits augmented with heat and moisture. *Malar J* 9: 6.
97. Knols BG, Jong R De, Takken W (1995) Differential attractiveness of isolated humans to mosquitoes in Tanzania. *Trans R Soc Trop Med Hyg* 89: 604-606.
98. Takken W, Knols BGJ (2010) Olfaction in vector-host interactions 2.
99. Logan JG, Birkett MA, Clark SJ, Powers S, Seal NJ, et al. (2008) Identification of human-derived volatile chemicals that interfere with attraction of Aedes aegypti Mosquitoes. *J Chem Ecol* 34: 308-322.
100. Meijerink J, Braks MAH, Brack AA, Adam W, Dekker T, et al. (2000) Identification of Olfactory Stimulants for Anopheles gambiae from Human Sweat Samples. *Journal of Chemical Ecology* 26: 1367-1382.
101. Bernier UR, Kline DL, Barnard DR, Schreck CE, Yost RA (2000) Analysis of human skin emanations by gas chromatography/mass spectrometry. 2. Identification of volatile compounds that are candidate attractants for the yellow fever mosquito (Aedes aegypti). *Anal Chem* 72: 747-756.
102. James AG, Hyliands D, Johnston H (2004) Generation of volatile fatty acids by axillary bacteria. *Int J Cosmet Sci* 26: 149-156.
103. Verhulst NO, Andriessen R, Groenhagen U, Kiss GB, Schulz S, et al. (2010) Differential attraction of malaria mosquitoes to volatile blends produced by human skin bacteria. *PLoS One* 5: e15829.
104. Yilmaz B, Portugal S, Tran TM, Gozzelino R, Ramos S, et al. (2014) Gut microbiota elicits a protective immune response against malaria transmission. *Cell* 159: 1277-1289.
105. Galili U, Mandrell RE, Hamadeh RM, Shohet SB, Griffiss JM (1988) Interaction between human natural anti-alpha-galactosyl immunoglobulin G and bacteria of the human flora. *Infect Immun* 56: 1730-1737.
106. Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, et al. (2015) Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 350: 1084-1089.
107. Ganai SC, Sanos SL, Kalfass C, Oberle K, Johner C, et al. (2012) Priming of natural killer cells by nonmucosal mononuclear phagocytes requires instructive signals from commensal microbiota. *Immunity* 37: 171-186.
108. Gao L, Song X, Wang J (2020) Gut microbiota is essential in PGRP-LA regulated immune protection against Plasmodium berghei infection. *Parasit Vectors* 13: 3.
109. Meister S, Agianian B, Turlure F, Relógio A, Morlais I, et al. (2009) Anopheles gambiae PGRPLC-Mediated defense against bacteria modulates infections with malaria parasites. *PLOS Pathog* 5: e1000542.
110. Guégan M, Minard G, Tran FH, Van VT, Dubost A, et al. (2018) Short-term impacts of anthropogenic stressors on Aedes albopictus mosquito vector microbiota. *FEMS Microbiol Ecol* 94.

111. Hughes GL, Dodson BL, Johnson RM, Murdock CC, Tsujimoto H, et al. (2014) Native microbiome impedes vertical transmission of Wolbachia in Anopheles mosquitoes. *Proc Natl Acad Sci USA* 111: 12498-12503.
112. Gendrin M, Rodgers FH, Yerbanga RS, Ouédraogo JB, Basáñez MG, et al. (2015) Antibiotics in ingested human blood affect the mosquito microbiota and capacity to transmit malaria. *Nat Commun* 6: 1-7.
113. Gendrin M, Yerbanga RS, Ouedraogo JB, Lefèvre T, Cohuet A, et al. (2016) Differential Effects of Azithromycin, Doxycycline, and Cotrimoxazole in ingested blood on the vectorial capacity of malaria mosquitoes. *Open Forum Infect Dis* 3: ofw074.
114. Ayoola OO, Adeyemo AA, Osinusi K (2015) Concurrent bacteraemia and malaria in febrile Nigerian infants. *Trop Doct* 35: 34-36.
115. Bygbjerg IC, Lanng C (1982) Septicaemia as a complication of Falciparum malaria. *Trans R Soc Trop Med Hyg* 76: 705.
116. Organization WH. Global vector control response 2017-2030.
117. Ferguson NM (2018) Challenges and opportunities in controlling mosquito-borne infections. *Nature* 559: 490-497.
118. Gao H, Cui C, Wang L, Jacobs-Lorena M, Wang S (2020) Mosquito microbiota and implications for disease control. *Trends Parasitol* 36: 98-111.
119. Abdul-Ghani R, Al-Mekhlafi AM, Alabsi MS (2012) Microbial control of malaria: Biological warfare against the parasite and its vector. *Acta Trop* 121: 71-84.
120. Cirimotich CM, Clayton AM, Dimopoulos G (2011) Low- and high-tech approaches to control Plasmodium parasite transmission by Anopheles mosquitoes. *J Trop Med* 2011: 891342.
121. Bawin T, Seye F, Boukraa S, Zimmer JY, Raharimalala FN, et al. (2016) Production of two entomopathogenic Aspergillus species and insecticidal activity against the mosquito Culex quinquefasciatus compared to Metarhizium anisopliae. 26: 617-629.
122. Seye F, Bawin T, Boukraa S, Zimmer JY, Ndiaye M, et al. (2014) Pathogenicity of Aspergillus clavatus produced in a fungal biofilm bioreactor toward Culex quinquefasciatus (Diptera: Culicidae). *J Pestic Sci* 39: 127-132.
123. Mohammed BR, Yayo AM, Deeni Y (2015) Insecticide resistance to Anopheles spp. Mosquitoes (Diptera: Culicidae) in Nigeria: A Review Malaria Project View project Role of NRF2 in ErbB targeted therapy View project. *Artic Int J Mosq Res*.