



ORIGINAL ARTICLE

Prevalence of Intestinal Parasitic Infections in Iraq during a Period from 2000-2020

Nadham Kadham Mahdi, M.Sc, Ph.D*

Department of Microbiology, College of Medicine, University of Basrah, Basrah, Iraq

*Corresponding author: Nadham Kadham Mahdi, M.Sc, Ph.D, Department of Microbiology, College of Medicine, University of Basrah, Basrah, Iraq



Abstract

The prevalence history for intestinal protozoal and helminthic infections among Iraqi people is reviewed from 2000-2020. The distribution of the intestinal parasites is mentioned in different communities including hospital patients, primary school children, food handlers, general population, immunocompromised patients, malnourished patients, sicklers and day-care centers attendance in both rural and urban regions. The prevalence rates are also illustrated in relation to age and sex parameters. Protozoal infections are the most common infections in Iraq. Other parasitic infections are of a local public health problem. Even parasitic infections could cause malabsorption, malnutrition and blood loss but also might cause physical and mental growth retardation specially among children. Therefore, Implantation of a national control program is essential to apply.

Keywords

Intestinal parasites, Iraq, Prevalence

Introduction

Intestinal parasites are prevalent throughout the world particularly in tropical and subtropical countries where deficiency in water and sanitation [1]. The high prevalence recorded in some regions possibly by introduction of infections through refugee camps and migration due to war and insecurity [2]. Many investigations have shown various prevalence rates of infections. The distribution of the parasites depends on the presence of suitable hosts, habits and favorable environmental conditions. Intestinal protozoa require only slight carelessness in habits leading to cosmopolitan distribution all over the world. They have a major public health importance due to their relation

to socioeconomic, environmental factors, sanitation level and education in the community. Intestinal helminthiasis contribute to poor nutritional status in infected children due to their adverse effects on food intake, normal nutrient absorption and immunity [3-5].

Nutritional impairment is often associated with chronic helminthiasis, as with those suffering from protein-energy malnutrition, iron-deficiency anemia and vitamin A deficiency [6,7]. The pathogenic protozoa are a common cause of diarrhea and have a worldwide distribution [6-8]. This remain an important cause of deaths among children less than 5 years old [9]. This is due to poorly developed immune systems. Therefore, severity of infection depends on virulence of the strain, host susceptibility and immune status as well as mixed infections [10].

Estimation of the global prevalence of the soil-transmitted helminths; 1000 million cases for *Ascaris lumbricoides*, 900 million for hookworms and 500 million for *Trichuris trichiura*. It has been reported that, probably 480 million people carried *Entamoeba histolytica* and 36 million develop invasive forms of amoebiasis. In addition, 200 million occur per year in Africa, Asia and Latin America [7,11].

In Iraq, studies on intestinal parasites started as early as 1939 [12]. Then, surveys were conducted recently and involved rural areas, school children in both sexes but mainly in and around the capital Baghdad. The prevalence diseases in Iraq differ from one region to another.

The present work illustrates the prevalence of intestinal parasites in relation to age, sex and socioeconomic factors.



Citation: Mahdi NK (2022) Prevalence of Intestinal Parasitic Infections in Iraq during a Period from 2000-2020. J Infect Dis Epidemiol 8:268. doi.org/10.23937/2474-3658/1510268

Accepted: June 29, 2022; Published: July 01, 2022

Copyright: © 2022 Mahdi NK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1: Map for Iraqi provinces.

Results and Discussion

The prevalence rates of intestinal protozoa and helminths parasites are reviewed according to the types of communities including general and selected populations in different Iraqi provinces (Figure 1).

Hospital patients

In and out patients are easier to investigate for parasitic infections. Ten surveys among all age groups were done in the middle part of the country, 4 in the Southern part and 7 in the Northern part (Table 1). However, medical history, morbidity and mortality of various parasitic infections are lacking in Iraq. Prevalence rates ranged from 7.36% to 84.67% and 0% to 18.01% of the protozoa and helminth infections respectively.

School children

School children (6-12 years old) were chosen by many workers because they are easier to deal with, represent the actual population of the communities and in various standards of living (Table 2). These data are concerned with the prevalence rates of infections among school children and not the household members of infected children.

Animal-handlers

Animal handlers including veterinarian, butchers and breeders could also acquire the parasitic infection at a rate of 50% in comparison to 14.8% for non-animal handlers (Table 3).

General population

The prevalence of intestinal protozoa and helminths among general populations in the middle part of the country including the capital Baghdad is illustrated in Table 4. The large samples surveyed were fairly representative of the general population. Even the prevalence rates of intestinal protozoa and helminths recorded during 2000-2020, were markedly lower than those reported in 1939 (73.9 and 41.0 for protozoal and helminths respectively) [12] but they are still representing a significant level from the public health point of view. Even so, it implants an improvement in sanitation, health and medical services.

Immunocompromised patients

Higher prevalence rates were recorded among patients with malignant diseases due to immunosuppression caused by either the disease itself

Table 1: The prevalence of intestinal protozoa and helminths among hospital's patients.

| Reference | No. Examined | Patients | % Infected | |
|-----------------------------|--------------|---|------------|----------|
| | | | Protozoa | Helminth |
| Yilmaz & Abdullah, [21] | 77 | Diarrhoeic patients from Duhok and Erbil. | 16.02 | 0 |
| Al-Joudi & Ghazal, [22]. | 6330 | Patients of 1-10 years old from Ramadi. Giardiasis is the most common infection. | | |
| | | During 1992-94. | 42.5 | 6.8 |
| | | During 1995-97. | 64.26 | 5.62 |
| Rhadi, et al. [23]. | 1026 | Patients 6-30 years old attending General Hospital in Basrah. <i>Cryptosporidium parvum</i> Recorded the highest infection (69) cases in age (20-30) years. | 46.44 | 1.7 |
| Al-Kubaisy, et al. [24]. | 2033 | Patients from Baghdad. Parasitic diarrhea was 22%. | 34.49 | 7.53 |
| Hussein & Meerkhl, [25]. | 1178 | Children attending Hivi Pediatric Hospital, Duhok Province. High infection in summer and rural area. Males were infected more than females. Highest rate among children more than 9 years. Artificially feeding had higher rate of infection. | 84.67 | 18.01 |
| AL-Khikani, et al. [26]. | 3176 | Patients with different ages attending the Al-Shomally General Hospital in Babil Province, ages 15-44 years are highly infected. Males and females are almost equal. | 11.0 | 0 |
| Hasan, et al. [27] | 615 | Patients attended Tikrit Teaching Hospital, 1-16 years old. The highest rate of infection in rural area and among age group 4-6 years. Males showed higher rate than females. | 14.3 | 0 |
| Salman, et al. [28] | 1071 | Children attending Paediatrics hospital in Al-Kadhimia, Al-Karkh side, Baghdad city, aged from seven days to 12 years. No difference between males and females. | 15.2 | 0.55 |
| Bazzaz, et al. [29] | 780 | They were admitted to Samarra General Hospital. The highest rate of infection was among 1-20 years old. | 8.35 | 0 |
| Murad, et al. [30] | 3967 | They were attended Azadi Teaching Hospital in Duhok City, Kurdistan Region. Males were more affected than females. | 14.7 | 0.07 |
| Al-Taei [16] | 2877 | Al-Hashimiyah hospitals, Babylon province/Iraq, among the ages from less than one year to 71 years old. The largest percent of infection reported in the (1-10) age group which was (44.35%). Male and female were 56.96 % and 43.03 %, respectively. | 49.4 | 0.2 |
| Al-Saad & Issa [31] | 428 | Children presenting at the pediatric hospital with a complaint of gastroenteritis. The age group 10-12 years showed the highest rate of infection (81.1%), and the lowest rate was in children 7-9 years (22.9%). | 31.3 | 0 |
| Al-Ibrahimi & Al-Waaly [32] | 109 | Surgical appendectomy in Women and Children Hospital in Al-Diwaniya showed <i>E. vermicularis</i> only. | -11.92 | 0 |
| Alomashi & Al-Shabbani [33] | 974 | Patients with anemia attending Al-Diwaniyah Teaching Hospital (Al-Qadisiyah province) | 64.0 | 64.0 |
| Kanabe & Darogha [17] | 548 | Children of both sexes 3 months to 10 years were attending Raparin Pediatric hospital in Erbil city | 7.36 | 0.18 |
| Mahdi & Ali [34] | 205 | Population aged 2 months -65 years old suffering from chronic diarrhoea for at least 2 weeks & attending the Maternity & Child Hospital in Basrah. 9.7% excreting <i>Cryptosporidium</i> oocysts. | 44.87 | 4.8 |
| Farhan [15] | 1804 | Males and females attending the Al Gailani-Central Medical Laboratory, Al-Ramadi/Al-Anbar, West of Iraq. | 38.25 | 3.35 |
| Husein, et al. [35] | 730 | People attending the Protecting Hospital for Pediatrics and Al-Zahrawy Hospital in Baghdad. | 90.1 | 17.5 |

| | | | | |
|-------------------------|-----|--|-------|------|
| Al-Hamairy, et al. [36] | 389 | Anemic patients of 1-50 years old were attended the primary health centre in Al-Doullab village-Babylon. | 31.4 | 9.29 |
| Al-Morshidy [37] | 243 | Patients attending the specialized Marjan Hospital for Internal and Cardiac Diseases in Hilla City-Babylon. Higher incidences were reported in patients using river water (80.8%) and low incidence among patients drinking tap water (43.8%). | 20.9 | 3.5 |
| Ali & Mohammed [38] | 307 | Children attending the Pediatric Teaching Hospital in Sulaimani city. | 22.15 | 0 |

Table 2: The prevalence of intestinal protozoa and helminths among primary school children.

| Reference | No. examined | Primary school children | % Infected | |
|-----------------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Al-Saad & Issa [31] | 451 | Children from primary school in Dohuk city. | 41.2 | 0 |
| Al-Ibrahimi & Al-Waaly [32] | 419 | Children of 1-12 years old of both sexes from schools and kindergartens in Al- Al Diwaniya. They reported <i>E. vermicularis</i> only. | - 43.67 | |

Table 3: The prevalence of intestinal protozoa and helminths among animal handlers.

| Reference | No. examined | Animal handlers | % Infected | |
|------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Mahdi & Ali [39] | 60 | Animal handlers (veterinarian, butchers & breeders) with mean age 27.16 ± 16.86 years. | 20.0 | 9.0 |

Table 4: The prevalence of intestinal protozoa and helminths among general population.

| Reference | No. examined | General population | % Infected | |
|----------------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Saheb [40] | - | Males and females from Baghdad, Karkh region. | 16.46 | 0 |
| Salman, et al. [41] | 417 | Displaced people in Kirkuk. Age 1-60 years. Age 1-10 years highly infected by <i>G. lamblia</i> . Gender was insignificant. | 3.42 | 0.23 |
| Salman, et al. [42] | 780 | Displaced people in Kirkuk. Age 1-78 years. Gender was insignificant. | 37.05 | 0.51 |
| Khudhair [43] | 5258 | Among Residents of Hawler, Soran and Chamchamal Cities, North of Iraq. Males were infected more than females. | 4.2 | 0 |
| Al-Shirifi & Abdullah [44] | 325 | Food handlers and food sellers in Kirkuk city. The highest infection rate among restaurant workers (12%). | 8.0 | 2.5 |
| Al-Naemy, et al. [45] | 422 | Children of Bashika District, suburb of Mosul city and 12 Km northeast of it. Males are affected more than females. | 32.6 | 5.7 |
| Mero & Hussein [46] | 1132 | Children of various sexes and ages in Dohuk and nearby villages. | 27.1 | 26.1 |
| Al-Daody, et al. [47] | 4062 | People of Erbil city. Females were infected more than males. | 0.23 | 4.04 |
| Jarallah [13] | 294 | children (159 males and 135 females) from 1 < to 14 years old in two rural villages (Abu-Malah and Harer) in Basrah marshlands regions | 27.4 | 17.87 |

or the cytotoxic drugs or both of them (Table 5).

Malnutrition

Only one work has been done in Basrah on 2005 where the malnourished children with diarrhea showed a higher prevalence rate (15.4%) for protozoal infections in comparison to malnourished children without diarrhea (10.75%) for protozoa and 2% for helminths (Table 6).

Sickle-cell anemia

Higher prevalence rate of parasitic infection (62.5%) was observed among patients with sickle-cell anemia in comparison with non-sicklers (14.8%) (Table 7).

Day-care centre

Almost similar prevalence rates were reported in

Table 5: The prevalence of intestinal protozoa and helminths among immunocompromised patients.

| Reference | No. examined | Immunocompromised patients | % Infected | |
|-------------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Mahdi, et al. [48] | 101 | 62 Males and 39 females with malignant diseases attending Oncology Centre of Basrah Teaching Hospital. Age 9 months-16 years old. | 28.7 | 10.9 |
| Mahdi & Al-Saadoon [49] | 58 | Males (37) and females (21) attending the Oncology centre of Basrah Maternity & Child Hospital. Microsporidium was at a rate 10.34% among Hodgkin and non-hodgkin lymphoma (83.3%) | 32.75 | 0 |

Table 6: The prevalence of intestinal protozoa and helminths among malnourished people.

| Reference | No. examined | Malnutrition | % Infected | |
|--------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Mahdi, et al. [50] | 194 | Malnourished children < 5 years old attending Basrah Maternity & Child Hospital. | | |
| | | Malnutrition with diarrhea. | 15.4 | 0 |
| | | Malnutrition without diarrhea. | 10.75 | 2.0 |

Table 7: The prevalence of intestinal protozoa and helminth among patients with sickle-cell anemia.

| Reference | No. examined | Sickle -cell anemia | % Infected | |
|------------------|--------------|--|------------|----------|
| | | | Protozoa | Helminth |
| Mahdi & Ali [51] | 40 | Stool samples were obtained from individuals admitted to three hospitals in Basrah with sickle-cell anemia. The first report for isolation of <i>Isospora belli</i> from a sickle-cell patient in Iraq and the Mediterranean region. | 55.0 | 7.5 |

Table 8: The prevalence of intestinal protozoa and helminths among the attenders of day-care center.

| Reference | No. examined | Day-Care Centre | % Infected | |
|---------------------|--------------|---|------------|----------|
| | | | Protozoa | Helminth |
| Mahdi & Ali [52] | 40 | 31 (72%) children were found to be infected for intestinal parasites compared to 1 (10%) positive personnel and 19 (34.5%) positive household contacts of the children. <i>Cryptosporidium</i> oocysts were found to be excreted in 4 (9%) children compared to 4 (7.2%) household contacts. No single positive case was recovered among the examined personnel of the day-care centre. | 42.5 | 27.5 |
| Al-Saad & Isaa [31] | 382 | Boys and girls attending the day-care center in Dohuk. | 43.3 | 0 |

Basrah (42.5%) and Dohuk (43.3%) as far as protozoal infections are concerned (Table 8).

Rural districts

Prevalence of intestinal parasitic infections is expected to be higher in rural districts and villages especially among children (Table 9). This can be attributed to the lower standard of hygiene, overcrowding, sanitation and probably food and/or water-borne infection. Agricultural practices and animal breeding are also responsible for the recorded prevalence rates. In marsh region of South Basrah 36.4% of water samples were contaminated with intestinal parasites [13]. Nevertheless, due to the progress in establishing and developing a good water supply,

electricity, paved roads and education, the prevalence rates were dropped in 2002 for helminthiasis but not for protozoan infections.

Urban districts

In contrast to the rural districts, the prevalence rates of intestinal parasites are markedly lower (Table 9). The recorded low prevalence rates are probably attributed to the adequate standards of living and conditions of the environmental sanitation.

Distribution according to sex

No significant differences were noticed in the prevalence rates of males and females (Table 9). But nevertheless, some works have reported the

Table 9: Distribution of intestinal parasites according to district and sex.

| Reference | No. examined | Province | Locality | | Sex | |
|-------------------------|--------------|----------------|----------|-------|-------|--------|
| | | | Urban | Rural | Male | Female |
| Hasan, et al. [27] | 615 | Tikrit | 11.54 | 20.87 | 17.3 | 11.22 |
| Bazzaz, et al. [14] | 780 | Samarra | - | - | 46.3 | 53.6 |
| Al-Taei [16] | 2877 | Al-Hashimiyah | - | - | 56.96 | 43.03 |
| Husein & Meerhkan [25] | 1172 | Dohuk | 34.09 | 65.9 | 63.98 | 36.02 |
| AL-Khikani, et al. [26] | 3176 | Babil | - | - | 50.65 | 49.35 |
| Khaudhair [43] | 5258 | Hawler | - | - | 4.6 | 3.4 |
| Kanabe & Drogha [17] | 548 | Erbil | - | - | 13.68 | 7.05 |
| Hamad & Ramzy [53] | 200 | Erbil | 34.69 | 25.49 | 27.35 | 33.73 |
| Al-Nawmy, et al. [45] | 422 | Bashika, Mosul | - | - | 60.6 | 39.33 |
| Salman, et al. [41] | 417 | Kirkuk | - | - | 4.55 | 5.75 |
| Farhan [15] | 1804 | Al-Anbar | - | - | 19.1 | 20.7 |
| Hussein, et al. [35] | 730 | Baghdad | - | - | 59.9 | 56.2 |
| Mero & Hussein [46] | - | Dohuk | 22.9 | 28.7 | 29.6 | 22.1 |
| Al-Hamairy, et al. [36] | 389 | Babylon | -60.15 | - | 60.9 | 58.7 |
| Ali & Mohammed [38] | 307 | Sulamini | 22.64 | 20.55 | 20.9 | 23.8 |

Table 10: Distribution of intestinal parasites according to age.

| Reference | 0-10 | Nov-20 | 21-30 | 31-40 | 41-50 | 51-60 |
|-------------------------|-------|--------|-------|-------|-------|-------|
| Salman, et al. [41] | 15.15 | 6.97 | 9.33 | 6.12 | 4.54 | 6.97 |
| Farhan [15] | 31.25 | 18.1 | 10.1 | 17 | 7.7 | 5.3 |
| Hussein, et al. [35] | 51.5 | 55.1 | - | - | - | - |
| Al-Hamairy, et al. [36] | 83 | 91 | 25 | 25 | 10 | - |
| Ali & Mohammed [38] | 24.05 | 48.84 | - | - | - | - |
| Hamad & Ramzy [53] | 37.06 | 50 | - | - | - | - |

prevalence rate of infection was higher among females in comparison to males. This may be due to fact that these females are housewives responsible for a house work, agricultural practices and animal breeding [14].

Distribution according to age

Lack of published papers was observed in relation to the distribution of infection among people with an age more than 21 years (Table 10). This is possibly due to sampling difficulty for the population with this wide range of ages.

Prevalence of intestinal parasites

The presented data presented in Table 11. The protozoan parasites are highly prevalent in all examined groups including general population, hospital patients, school children in urban and rural districts reaching a rates of infection of 61.6 [15] and 88% [16]. The highest rate of infection reported for the emerging opportunistic *Cryptosporidium* was 18.94% [17]. A significant association was found between giardiasis and water contamination. About 31.6% of the inhabitants use storage water, while only 1.3% use sterile water bottles [18]. In addition, the use of waste water to irrigate vegetables or organic manure as a fertilizer that are eaten raw without proper washing leads to

food-borne parasitic infections [19,20]. Ascariasis was detected at a lower rates of infection (Table 11). In contrast, prevalence of hymenolepiasis nana was getting higher in 15 studies out of 21 indicating a public health problem specially among children. Thus, it needs more attention for application of control measure. *Enterobius vermicularis* was detected only accidentally because stool examination is of no value in its specific diagnosis [7].

Some workers have reached a decision that the concentration stool examination methods are better than direct smear method in the diagnosis of helminthiasis which might have an influence on the prevalence rate in the area or the examined population. Therefore, if this is true, there will be a close relationship between the rate of infection and the method followed for stool examination.

Conclusion

Although the prevalence rates of intestinal parasitic infections are varied from one area to another according to the degree of personal and community hygiene, sanitation and climatic factors are relatively common in Iraq. Intestinal protozoan, helminthic or mixed infections are prevalent among all types of communities and population samples from both urban and rural

Table 11: Distribution of intestinal protozoa and common helminths in Iraq.

| Reference | <i>E. histolytica</i> | <i>G. lamblia</i> | <i>B. hominis</i> | <i>A. lumbricoides</i> | <i>E. vermicularis</i> |
|------------------------------|-----------------------|-------------------|-------------------|------------------------|------------------------|
| Salman, et al. [28] | 27 | 11 | 7.6 | - | 0.3 |
| Bazzaz, et al. [14] | 12.8 | 3.9 | - | - | - |
| Al-Taei [16] | 88 | 10.8 | - | - | 0.03 |
| Yilmaz & Abdullah [21] | 24.45 | 7.6 | - | - | - |
| Rhadi, et al. [23] | 15.3 | 5.94 | 7.01 | 16.27 Crypto | 3.21 |
| Al-Kubaisy, et al. [24] | 23.44 | - | - | 2.2 | 12.7 |
| Hussein and Meerkhan [25] | 38.7 | 25.67 | 4.6 | 9.2 Crypto | 18.01 |
| Salman, et al. [41] | 1.67 | 9.35 | 4.17 | 1.43 Crypto | - |
| Al-Khikaniet al. [26] | 17.91 | 4.09 | - | - | - |
| Kanabe and Darogha [17] | 0.18 | 9.72 | 9.72 | 18.94 Crypto | - |
| Al-Daoody, et al. [47] | 0.12 | 0.34 | - | - | - |
| Al-Shirifi and Abdullah [44] | 3.69 | 12.31 | - | 1.85 | 3.38 |
| Al-Nawmy, et al. [45] | 28.2 | 37 | - | 3.8 | 15.3 |
| Al-Bazzaz, et al. [14] | 12.8 | 3.9 | - | - | - |
| Al-Taei [16] | 12.14 | 4.9 | - | 0.37 | 0.38 |
| Farhan [15] | 61.6 | 14.9 | - | 6.7 | 9.8 |
| Hussein, et al. [35] | 24.3 | 31 | - | 9.7 | - |
| Mero and Hussein [46] | 10.15 | 11.92 | 0.08 | - | 0.35 |
| Al-Morshidy [37] | 27.9 | 13.9 | - | 2.8 | 2.4 |
| Al-Hamairy, et al. [36] | 36.7 | 26.1 | - | 1.3 | 32.9 |
| Jarallah [13] | 23.87 | 30.93 | - | 2.58 | 47.46 |
| Mahdi & Ali [39] | 7.5 | 28 | 36 | 5.0 Crypto | 2.5 |

Crypto = *Cryptosporidium* species.

regions of Iraq. Adults and children of both sexes are infected. Therefore, an urgent and efficient preventive and control measures is essential.

Even an adequate treatment for amoebiasis and giardiasis, which are the common causes for diarrhea, is available, the morbidity is still high. The complications of invasive amoebiasis are potentially fatal and giardiasis may cause malabsorption in children [4,6,7].

Several intestinal helminthiasis contribute to the general and persistence of malnutrition and reduced productivity [6,7]. Careful evaluation of the individual parasite should be continued in different regions and among different societies.

Implementing a national control program should include a primary health care, health education, family planning, water supply, paved roads, environmental sanitation, nutrition, mass treatment by new broad spectrum anthelmintics, avoid crowding and improvement in agriculture practices. Anti-protozoal and broad-spectrum anthelmintics are available and effective at a regular mass treatment. Well trained health workers chosen from the same community are valuable in the diagnosis and treatment of various parasitic infections especially in rural areas and far villages in the country.

References

- Damen JG, Luka J, Biwan EI, Lugos M (2011) Prevalence of intestinal parasites among pupils in Rural North Eastern, Nigeria. *Niger Med J* 52: 4-6.
- Rashid MK, Joshi M, Joshi HS, Fatemi K (2011) Prevalence of intestinal parasites among school going children in Bareilly District. *NJIRM* 2: 35-36.
- Crompton DWT, Nesheim MC, Pawlowski ZS (1985) Ascariasis and its public health significance. Taylor and Francis, Basingstoke.
- Pereira PCM (2003) Interaction between infection, nutrition and immunity in tropical medicine. *J Venom Anim Toxins incl Trop Dis* 9: 163-173.
- Shea-Donohue T, Qin B, Smith A (2017) Parasites, nutrition, immune responses, and biology of metabolic tissues. *Parasite Immunol* 39: 10.
- WHO Expert Committee (1987) *Bulletin of the World Health Organization* 65: 575.
- John DT, Petri WA (2006) *Markell and Voge Medical Parasitology*. Saunders Elsevier Inc.
- Weber R (2020) *Hunter's tropical medicine and emerging infectious diseases*.
- United Nations International Children's Emergency Fund (1990) *Annual report of communicable disease report control centre, Iraq*.
- Stanley SJ (2003) Amoebiasis. *Lancet* 361: 1025-1034.
- Frias L, Leles D, Araujo A (2013) Studies on protozoa in ancient remains - A review. *Mem Inst Oswaldo Cruz* 108: 1-12.

12. Senekji HA, Boswell C, Beattie CP (1939) Incidence of intestinal parasites in Iraq. *Transaction of the Royal Society Tropical Medicine and Hygiene* 33: 349-352.
13. Jarallah HM (2012) Intestinal parasitic infections among rural villages in Basrah marshes regions. *Journal of Basrah Researches (Sciences)* 38: 40-43.
14. Bazzaz AA, Ouhood M, Shakir OM, Alabbasy RH (2017) Prevalence of two gastrointestinal parasites *Entamoebahistolytica* and *Giardia lamblia* within Samarra City, Iraq. *Advances in Bioscience and Biotechnology* 8: 399-410.
15. Farhan AO (2012) Prevalence of Intestinal parasitic infestations in Al-Anbar province, West of Iraq. *J University of Anbar for Pure Science* 6: 1991-1995.
16. Al-Taei AHO (2019) The prevalence of intestinal parasite among the attending people to Al-Hashimiyah hospitals for seven years, Babylon province, Iraq. *Journal of Physics: Conference Series* 1294: 062022.
17. Kanabe LO, Darogha SNR (2017) Epidemiology of Cryptosporidiosis among diarrheic children of Raparin Pediatric hospital, Erbil province-Kurdistan Region, Iraq. *Cihan University-Erbil Scientific Journal* 2: 538.
18. Muslim F (2014) The prevalence of *Giardia lamblia* among infants and young children in Wasit governorate, Iraq. *Science Medicine* 7: 181-189.
19. Damen JG, Banwat EB, Egah DZ (2007) Parasitic contamination of vegetables in Jos, Nigeria. *Ann Afr Med* 6: 115-118.
20. Mohamed MA, Siddig EE, Elaagip AH, Edris AMM, Nasr AA (2016) Parasitic contamination of fresh vegetables sold at central markets in Khartoum state, Sudan. *Ann Clin Microbiol Antimicrobio* 15: 17.
21. Yilmaz H, Abdullah AM (2017) Prevalence of intestinal parasites (*Entamoeba* species and *Giardia lamblia*) in Duhok and Erbil cities, Northern Iraq. *J Microbiol Exp* 4: 119-122.
22. Al-Joudi FS, Ghazal AM (2005) The prevalence of intestinal parasites in Ramadi, Iraq. *Bulletin of Pharmaceutical Science. Assiut* 28: 277-291.
23. Rhadi HA, Abdul-Zahra A, Abdul-Jabar S (2018) Prevalence of intestinal pathogenic parasites in Basrah City. *Int J Sci Res* 8: 174-180.
24. AL-Kubaisy W, AL-Talib H, Al-khateeb A, Shanshal MM (2014) Intestinal parasitic diarrhea among children in Baghdad-Iraq. *Trop Biomed* 31: 499-506.
25. Hussein JN, Meer Khan AA (2019) The incidence of intestinal parasites among children in Hivi pediatric hospital, Duhok, Iraq. *Science Journal of University of Zakho* 7: 1-4.
26. AL-Khikani FH, Almosawey HA, Hameed RM, Alhussain BA, Ayit AS, et al. (2019) Prevalence of *Entamoebahistolytica* and *Giardia lamblia* associated with infectious Diarrhea in Al-Shomally population, Babil, Iraq. *Biomed Biotechnol Res J* 3: 245-248.
27. Hasan TAH, Muhaimid AA, Mahmoud AR (2020) Epidemiological study of *Giardia lamblia* in Tikrit city, Iraq. *Systemic Review in Pharmacy* 11: 102-106.
28. Salman AO, Mhaisen FT, Abdul-Rahman A, Al-Tae A (2019) On the occurrence of some intestinal parasites among Diarrheic children attending a hospital in Al-Karkh Side, Baghdad City, Iraq. *Biological and applied Environmental Research* 3: 93-102.
29. Bazzaz AA, Hamad HR, Shakir OM, Bazzaz SA (2019) Evaluation of two gastrointestinal parasites *Entamoebahistolytica* and *Giardia lamblia* within Iraq. *J Community Med Health Educ* 8: 21-22.
30. Murad AM, Al-Saad ATM, Mustafa AA (2018) Prevalence of intestinal parasites among patients attending azadi teaching hospital in Duhok city-Kurdistan region/Iraq. *Pure and Engineering* 21: 68-74.
31. Al-Saeed A, Issa S (2006) Frequency of *Giardia lamblia* among children in Dohuk, northern Iraq. *East Mediterr Health J* 12: 555-562.
32. Al-Ibrahimi MHM, Al-Waaly ABM (2020) The prevalence of pinworm, incontinence, and appendicitis in children in Diwaniyah governorate, Iraq. *Annals of Tropical Medicine and Public Health* 23: 5509.
33. Alomashi GBA, Al-Shabbani AH (2019) Prevalence of intestinal parasitic infestation in anemic patients attended to Al-Diwaniyah Teaching Hospital at Al-Qadisiyah Province/Iraq. *International Journal of Pharmaceutical Quality Assurance* 10: 60-65.
34. Mahdi NK, Ali NH (2004) Cryptosporidiosis and other intestinal parasitic infections in patients with chronic diarrhea. *Saudi Med J* 25: 1204-1207.
35. Hussein RA, Shaker MJ, Majeed AA (2011) Prevalence of intestinal parasitic infections among children in Baghdad City. *Journal of the College of Basic Education* 17: 21-30.
36. Al-HamairyAKA, Al-Mosauy AMA, Al-Rubaea AHM (2013) Prevalence of parasitic infection and relationship with anemia in Al-Doullab village, Babylon province, Iraq. *Egypt J Exp Biol (Zool)* 9: 231-236.
37. Al-Morshidy M (2006) Prevalence of intestinal pathogenic parasites infections in Hilla city, Babylon province, Iraq. *Journal Babylon University of Pure Applied Science* 14: 179-185.
38. Ali SA, Mohammed LO (2010) Prevalence of intestinal parasite among children in Sulaimani city. *Journal of Duhok University* 13: 94-98.
39. Mahdi NK, Ali NH (2002) Cryptosporidiosis among animal handlers and their livestock in Basrah, Iraq. *East Afr Med J* 79: 550-553.
40. Saheb EJ, Mahdi SG, Mosa IS, AbdulKarim MI, Khistawi AN (2017) Epidemiology of some parasitic helminthes in Iraq from 2011 until 2015. *Iraqi Journal of Science* 58: 789-796.
41. Salman YJ, Al-Tae AA, Abid AM (2016) Prevalence of *Giardia lamblia* among Iraqi displaced peoples in Kirkuk province. *Int J Curr Microbiol App Sci* 5: 753-760.
42. Salman YJ, Sadek WS, Rasheed ZK (2015) Prevalence of *Cryptosporidium parvum* among Iraqi displaced people in Kirkuk city using direct microscopy, flotation technique and ELISA-copro antigen test. *Int J Curr Microbiol App Sci* 4: 559-572.
43. Khudhair A (2020) Prevalence of *Giardia lamblia* among Residents of Hawler, Soran and Chamchamal Cities, North of Iraq. *Pak-Euro Journal of Medical and Life Sciences* 3: 28-36.
44. Al-Shirifi HM, Abdullah IA (2005) Prevalence of intestinal parasites amongst food handlers in Kirkuk city, Al-Taam'ern province, Iraq. *Rafidain Journal of Science* 16: 13-21.
45. Al-Naemy BS, Al-Kalak S, Rahemo ZIF (2012) The intestinal parasites of Bashiqa District, Nineveh Governorate, Iraq. *Int J Molecular Zoology* 2: 51-54.

46. Mero WMS, Hussein JN (2013) Prevalence of intestinal parasites among children in various localities of Duhok city and nearby villages. *Science Journal of University of Zakho* 1: 189-199.
47. Al-Daody AAK, Younes MR, Ali WR, Hamad KM (2017) Prevalence of *Hymenolepis nana* in Erbil City - North of Iraq. *International Journal of Research Studies in Science, Engineering and Technology* 4: 16-21.
48. Mahdi NK, Al-Sadoon MA, Hassan GK (2007) Cryptosporidiosis and immunological status in children with malignant diseases. *The Medical Journal of Basrah University* 25: 1-6.
49. Mahdi NK, Al-Sadoon MA (2012) Microsporidiosis among children with malignant diseases in Basrah, Iraq. *Pakistan Journal of Medical Sciences* 28: 621-624.
50. Mahdi NK, Jassim RM, Hassan MK (2005) Intestinal parasitic infections including cryptosporidiosis and immunological aspects among malnourished children. *Journal of the Bahrain Medical Society* 17: 43-48.
51. Mahdi NK, Ali NH (2002) Intestinal parasites including *Cryptosporidium* species in Iraqi patients with sickle cell anaemia. *East Mediterr Health J* 8: 345-349.
52. Mahdi NK, Ali NH (2002) Intestinal parasitic (including *Cryptosporidium*) infections in day-care centers. *Bahrain Medical Bulletin* 24: 135-137.
53. Hamad NR, Ramzy IA (2012) Epidemiology of *Entamoeba histolytica* among children in Erbil, Province, Kurdistan Region-Iraq. *Journal of Research in Biology* 1: 57-62.